

Research Article

Open Access

Solar Alpha Particles with High Energy Influence Neoplasms Mortality in Europe and the Mediterranean?

Nikolay Takuchev

Ph.D., Associate Professor, Trakia University, Stara Zagora, Bulgaria

ABSTRACT

A dangerous phenomenon for humanity is described. Through a joint analysis of data from satellites in orbit around the Earth and from the health statistics source EUROSTAT, it became clear that, by some mechanism, the influx of positively charged particles with high energy into the Earth's orbit increases mortality on the planet's surface. The increase in mortality is in a band of maximum risk in the Northern Hemisphere, parallel to the Equator and bounded by the parallels of 30° and 50° north latitude. Examples are given for the European Union mortality from neoplasms, confirming the described phenomenon. A hypothetical mechanism based on observational evidence has been proposed, according to which this dangerous phenomenon is caused by high-energy solar alpha particles that are sufficient to overcome the atmosphere's resistance and reach the Earth's surface in a limited area, resulting in a maximum death impact.

*Corresponding author

Nikolay Takuchev, Ph.D., Associate Professor, Trakia University, Stara Zagora, Bulgaria.

Received: July 19, 2025; **Accepted:** July 23, 2025; **Published:** July 31, 2025

Keywords: Mortality, Neoplasms, Satellites Goes, Cosmic Alpha Radiation, Eurostat

Introduction

In a series of publications, a potentially hazardous phenomenon of cosmic origin affecting humanity was described - the existence of a high correlation between the flows of positively charged particles with high kinetic energy recorded from satellites in Earth orbit and mortality on the Earth's surface from various diseases [1-25]. The main focus of the mentioned publications was on the effect of solar alpha radiation on mortality from diseases, the leading cause of death in the circulatory system. The harmful effects of solar alpha radiation are also observed in many other organs and systems of the human organism, rendering this invisible effect one of the primary causes of death for humanity. The mentioned dangerous phenomenon of cosmic origin is unevenly spread on the planet's surface. Europe and the Mediterranean are among the most affected. However, numerous examples are described in where this phenomenon is also observable in several countries from the Northern Hemisphere, including Asia, America, and even Africa [1-25]. The European Union has a tradition of maintaining reliable mortality statistics, in which the discussed multifaceted influence of solar alpha radiation on mortality stands out clearly. Below, based on mortality statistics from the European Union, the unexpected association between solar alpha radiation and mortality from neoplasms is illustrated. High-energy alpha radiation can ionize the atoms in living organisms, significantly impairing normal cellular metabolism. This type of radiation may be the leading external cause of neoplasms.

Materials and Methods

Mortality Data

The analysis below is based on an authoritative source of health data -EUROSTAT [26]. In the study, the parameter annual mortality rate – number of deaths per 100,000 inhabitants was used as a

characteristic of mortality. EUROSTAT offers free access to data on mortality rates from specific causes in the countries of the European Union, the European Economic Area, and the candidate countries for EU membership. Geographically, these countries are situated in Europe and the Mediterranean region. Data are grouped by NUTS (Nomenclature Des Unites Territoriales Statistiques in French, the nomenclature of territorial units for statistics). In the study, mortality data from the EUROSTAT shortlist were used, which categorizes mortality rates by cause of death into 92 groups, primarily diseases. The groups are related to the classes in the International Classification of Diseases (ICD-10), 10th revision. The shortlist includes mortality data for EU countries (NUTS-1) and EU regions (NUTS-2, which are smaller areas within the larger NUTS-1 countries). Currently (2023), the shortlist includes mortality rate data for the period 2011–2020. Annual mortality rate data were extracted for 353 European regions (NUTS-2) separately for each of the shortlisted groups for the period 2011–2019 (the last pre-pandemic year).

Satellite Data

Satellite data on corpuscular radiation-protons and alpha particles recorded by the satellites of the series GOES (Geostationary Operational Environmental Satellites) were obtained from a NOAA site [27].

The satellites of the GOES series fly in geostationary orbit (above the Earth's equator), at an altitude of 36,000 kilometers above the Earth's surface, make one lap in 24 hours, that is, they "hang" over a certain point on the Earth's surface and are not shade by the Earth at their circumference around it.

Data on alpha-particle and proton fluxes (unit: (number of particles). $\text{cm}^{-2} \cdot \text{s}^{-1} \cdot \text{sr}^{-1} \cdot \text{MeV}^{-1}$)) with energies in the range 3.8-21.3 MeV were used. The fluxes were recorded by the satellite high-energy particle

detectors: 1. Energetic Particles Sensor (EPS), and 2. Energetic Proton, Electron, and Alpha Detector (EPEAD). The data are available averaged over a 5-minute interval, during which there are up to 25 reports of the instrument.

Data Processing

The correlation coefficients between the annual averaged alpha radiation flux and the annual mortality rate were calculated [28].

Maps were created showing (with black isolines) the distribution of the annual mortality for 353 NUTS-2 European regions from the EUROSTAT shortlist death causes: “Neoplasms” (EUROSTAT mortality shortlist number 8), “Malignant neoplasms (C00-C97)” (shortlist number 9), and “Other malignant neoplasms (remainder of C00-C97)” (shortlist number 31) for 2012, the year with the highest solar activity (and mortality) in the studied time interval 2011 – 2019. Below, the three groups of causes are collectively referred to as “neoplasms”. With red isolines, the maps display the distribution of the correlation coefficient across Europe and the Mediterranean between the annual mortality rate from neoplasms and the annually averaged alpha particle flux for the time interval 2011-2019. Data on the coordinates, latitude, and longitude [Google Earth] of the centroids of the NUTS-2 regions included in the study were used in the map. Mapping was performed with Golden Software Surfer 10. The kriging interpolation procedure was selected.

In mathematical statistics, the level of statistical significance is a parameter that indicates the degree of reliability of the calculated correlation coefficient [28]. The smaller the value of this parameter, the more reliably the correlation coefficient is established, i.e., the more reliably a cause-and-effect relationship is established between the annual flux of solar alpha radiation and mortality from neoplasms.

The correlation coefficient and the level of statistical significance are related. For the 9 years included in the study, a minimum correlation coefficient of 0.668 corresponds to a statistical significance level of 0.05 [28]. In scientific studies, a level of statistical significance of 0.05 or less is typically accepted as a criterion for the reliability of the correlation coefficient. The red isolines on the correlation coefficient distribution map enclose regions with statistically significant values of the correlation coefficients, up to a significance level of 0.05. Correlation coefficients with a significance level above 0.05 are of high reliability (the higher the number, the lower the significance level), i.e., the existence of a causal relationship between cosmic alpha radiation and mortality from the causes of neoplasms can be considered reliably established in the mentioned areas enclosed by red isolines of the correlation coefficient.

If there is a coincidence between some of the maxima for mortality rates and correlation coefficients, then in the region of these maxima, the impact of alpha radiation contributes noticeably to mortality from neoplasms.

To the extent that the hypothetical mechanism proposed below, which explains the observed phenomenon, assumes that charged particles of high energy pass through the atmosphere and reach the Earth's surface, the energy required for this process was calculated using databases and calculators PSTAR and ASTAR [29,30]. Geomagnetic field data were obtained from the INTERMAGNET site [31].

Results

The described dangerous phenomenon is observed in the form of dependence between the annual average flux of radiation from positively charged particles (protons and alpha particles) with high kinetic energy, recorded by satellites in orbit around the Earth, and the annual mortality rate in the statistics of several countries from all continents in the Northern Hemisphere. The countries in whose mortality statistics the phenomenon is observed are located in a band parallel to the equator with approximate boundaries along the parallels of 30° and 60° north latitudes. It is observed in the annual mortality statistics of small countries. It is not noticeable in the statistics of large countries in the same band. It can be inferred that the impact on the Earth's surface is short-lived and limited to a small area, comparable to that of a small country, but is masked in large country statistics because it does not affect the entire area of the large country simultaneously. This conclusion is confirmed for the USA, for which data on mortality in individual states are available [10].

The mortality statistics of the European Union are suitable for the study because they are based on statistical regions, which are smaller than a country but still large enough to include a statistically sufficient number of inhabitants.

This phenomenon would be expected to influence mortality in countries south of the Equator; however, the mortality statistics for these regions are scarce, unreliable, or absent, preventing reliable inferences about its influence in the Southern Hemisphere.

For particle energies ranging from 3.8 to 21.3 MeV, the year-averaged fluxes of protons and alpha particles are highly correlated, indicating that the studied phenomenon of lethality is noticeable in both the mean proton flux and the mean alpha particle flux data. For the reason explained below, only the average flux of high-energy alpha particles is included as the incident radiation in the next examples.

With black isolines, Figure 1 shows the distribution of the mortality rate for EUROSTAT shortlist death cause „Other malignant neoplasms (remainder of C00-C97)” for Europe and the Mediterranean, for 2012, the year of maximum solar activity (and mortality), from the studied interval. With red isolines in the figure, the distribution of the statistically significant correlation coefficient between solar alpha particle fluxes and the mortality rate for “Other malignant neoplasms (remainder of C00-C97)” is shown. The statistically significant influence of alpha radiation on mortality from “Other malignant neoplasms (remainder of C00-C97)” is limited to a band parallel to the Equator with approximate dimensions between 30° and 60° north latitude. The impact of alpha radiation on mortality in the mentioned band does not cover the entire band continuously, but rather occurs in limited spots, primarily affecting mountainous areas. A pronounced effect - a statistically significant coincidence between areas with increased annual “Other malignant neoplasms (remainder of C00-C97)” mortality rate and its correlation with annual alpha radiation flux is observed for two narrow bands around the 40th and 55th parallels.




Figure 1: Europe and the Mediterranean: the EUROSTAT mortality rate for 2012, classified in Shortlist classification as “Other malignant neoplasms (remainder of C00-C97)” (black isolines), and its statistically significant correlation with annual alpha particle flux (red isolines) for the studied interval, 2011-2019.

Figures 2, 3, 5, 6, 7, and 9 show the time dependence in the interval 2011-2019 of two numerical sequences:

- Of the recorded annual fluxes of alpha particles from satellites of the GOES series – 10, 11, 12, 13, and 14, and,
- Of the “Neoplasms”, “Malignant neoplasms (C00-C97)”, and “Other malignant neoplasms (remainder of C00-C97)” annual mortality rate for some of the NUTS-2 regions of the European Union.
- The high correlation between the two numerical sequences is evident, suggesting a causal relationship between them.




Figure 2: The highly statistically significant correlation between annual fluxes of cosmic alpha radiation and the “Other malignant neoplasms (remainder of C00-C97)” mortality rate for the EUROSTAT NUTS-2 region Schweiz/Suisse/Svizzera, Switzerland, indicates the presence of a causal relationship between the two phenomena.




Figure 3: The statistically significant correlation between annual fluxes of cosmic alpha radiation and the “Other malignant neoplasms (remainder of C00-C97)” mortality rate for the EUROSTAT NUTS-2 region of Orta Anadolu, Türkiye, indicates the presence of a causal relationship between the two phenomena.

Figure 4 shows the distribution of the mortality rate for EUROSTAT shortlist death cause “Malignant neoplasms (C00-C97)” for Europe and the Mediterranean, under the same conditions and designations as in Figure 1. Areas with increased statistically significant correlation are scattered throughout Europe. A pronounced effect – a statistically significant coincidence between areas with increased annual “Malignant neoplasms (C00-C97)” mortality rate and its correlation with annual alpha radiation flux is observed for the two narrow meridional bands between -5° and $+5^{\circ}$ and 10° and 20° .




Figure 4: Europe and the Mediterranean, the “Malignant neoplasms (C00-C97)” mortality rate for 2012 (black isolines), and its statistically significant correlation with annual alpha particle flux (red isolines) for the studied interval, 2011-2019.




Figure 5: The high statistically significant correlation between annual fluxes of cosmic alpha radiation and the “Malignant neoplasms (C00-C97)” mortality rate for the EUROSTAT NUTS-2 region, Steiermark, Austria, indicates the presence of a causal relationship between the two phenomena.




Figure 6: The high statistically significant correlation between annual fluxes of cosmic alpha radiation and the “Malignant neoplasms (C00-C97)” mortality rate for the EUROSTAT NUTS-2 region Alsace, France, indicates the presence of a causal relationship between the two phenomena.




Figure 7: The high statistically significant correlation between annual fluxes of cosmic alpha radiation and the “Malignant neoplasms (C00-C97)” mortality rate for the EUROSTAT NUTS-2 region Drenthe, Netherlands, indicates the presence of a causal relationship between the two phenomena.

Figure 8 shows the distribution of the mortality rate for EUROSTAT shortlist death cause “Neoplasms” for Europe and the Mediterranean, under the same conditions and designations as in Figure 1. The areas with statistically significant correlations are scattered throughout Europe, and their pattern is similar to the mix of patterns in Figures 1 and 4.




Figure 8: Europe and the Mediterranean, the “Neoplasms” mortality rate for 2012 (black isolines), and its statistically significant correlation with annual alpha particle flux (red isolines) for the studied interval of 2011-2019.




Figure 9: The statistically significant correlation between annual fluxes of cosmic alpha radiation and the “Neoplasms” mortality rate for the EUROSTAT NUTS-2 region, specifically West Central Scotland (NUTS 2021), United Kingdom, suggests the presence of a causal relationship between the two phenomena.

The link between diseases such as solar lentigines and type 2 diabetes mellitus, as well as malignant non-Hodgkin lymphoma and cutaneous melanoma, can also be explained by a common cause: the fluxes of high-energy solar alpha radiation [24,25].

Discussion

The given examples convincingly demonstrate the influence of cosmic radiation flux on neoplasm mortality. The mechanism of the described influence remains unclear. A hypothesized mechanism of this influence is outlined below, answering many of the questions that arise.

- 1. An Observed Phenomenon**-mortality from many diseases in the statistics of many countries located mainly in the 30°N-60°N band, is strongly correlated with fluxes of positively charged particles with energy of the order of 4-21 MeV, recorded by the GOES series satellites in Earth orbit.
- 2. Proposed Hypothesis**-positively charged particles with high energy penetrate through the Earth’s atmosphere to the Earth’s surface and damage human health, causing death mainly in elderly people.

3. Positively charged solar particles capable of penetrating through the Earth's atmosphere to the Earth's surface are high-energy alpha particles. Calculators PSTAR [28] and ASTAR [29] calculate the penetration parameters of protons, respectively alpha particles, in different substances, in particular in air. Calculations with data for a homogeneous atmosphere -an atmospheric model with constant density, temperature, and pressure decreasing with height show that only particles whose energy is above 2.4 GeV for protons and over 6.2 GeV for alpha particles can penetrate the Earth's atmosphere to the surface [4]. There are no protons registered by GOES satellites above 0.7 GeV; however, alpha particles with energies above 3.4 GeV are registered, and hypothetically, those with energies above 6.2 GeV are also detected [4,10, and 12]. Therefore, the particles that penetrate the Earth's surface are likely high-energy alpha particles. Only fluxes of alpha particles with a magnitude of at least $1000 \text{ particles} \cdot \text{cm}^{-2} \cdot \text{s}^{-1} \cdot \text{sr}^{-1} \cdot \text{MeV}^{-1}$ is correlated with the mortality of the Earth's surface. The recorded alpha particle flows are mostly pulses, with durations ranging from less than 5 minutes (the averaging interval of the recording device) to a few hours.
4. As the average altitude of the affected countries increases, the particle flux-correlated mortality shows an increasing trend [4,10]. It is likely due to the more intense radiation flux penetrating the thinner atmosphere over the mountainous regions of Earth's surface -an argument that favors the hypothesis.
5. The source of the flows of positively charged particles is the Sun – mortality increases with observable processes on the Sun-from Solar Mass Ejections directed to Earth (a phenomenon on the solar surface that could be observed with other astronomical means) [4,8]. The Alpha Magnetic Spectrometer (AMS-02) on the International Space Station measures cosmic rays, excluding those of solar origin (when shielded from the Sun by the station's solar panels). In particular, it measures the flow of ^3He and ^4He (alpha particles) in cosmic rays [32]. The measurements show increasing annual flux of alpha particles in cosmic rays for the interval of years from 2011 to 2017 (last available data), while the flux of GOES registered (solar?) alpha particles for the same interval of years is decreasing (Figures 2 and 9, for example). Indirect evidence for the Sun as a source of high-energy alpha particles is that this assumption convincingly explains the downstream processes that ultimately lead to deaths on Earth's surface.
6. It is assumed that the alpha particles recorded by the satellites were emitted simultaneously with the hypothetical fast alpha particles in a common explosive process (flares?) on the solar surface. It can be calculated that particles with an energy of 7 GeV require 8.87 minutes to reach the Earth's surface from the Sun's surface, while particles with energies of 5-10 MeV, as registered by the satellites, travel approximately 2 hours. The registered alpha particles do not have enough energy to penetrate the atmosphere, unlike the hypothetical fast alpha particles that reach the Earth's surface in minutes from the center of the Sun's disk. However, the registered alpha particles indicate that two hours earlier, the Earth's surface was irradiated with fast (unregistered) alpha particles.
7. Although alpha particle streams irradiate the entire illuminated part of the atmosphere, penetration of fast alpha particles to the surface occurs only in a limited area of the surface (death spot),

- a The Sun is culminating at the center of the death spot. During the year, the apparent position of the Sun relative to the point of observation shifts, so that the maximum angle of elevation of the Sun's disk above the horizon (solar culmination at local noon) changes depending on the date. The latitude and longitude of a point on the Earth's surface where the Sun's disk is at its culmination at that moment of registration of the incoming alpha particle flow-the point of registration, can be determined from the date (latitude), and the hours and minutes of registration (longitude). The center of the dead spot can be calculated-it is approximately 30° east of the registration point [10]. The Earth's angular velocity is 15° per hour. Figure 10 shows the registration points for cases of intense alpha radiation fluxes with a magnitude of at least $1000 \text{ particles} \cdot \text{cm}^{-2} \cdot \text{s}^{-1} \cdot \text{sr}^{-1} \cdot \text{MeV}^{-1}$ for 1974 and 1975. They are located on meridians approximately 30° west of the meridional bands of high correlation between alpha particle fluxes and mortality from "Malignant neoplasms (C00-C97)" (Figure 4). The strange meridional location of the points of registration of alpha radiation fluxes at the end of the 20th century is the subject of articles [10,12, 15, and 23].



Figure 10: Points of registration of pulses of alpha particles with a flux not less than $1000 \text{ particles} \cdot \text{cm}^{-2} \cdot \text{s}^{-1} \cdot \text{sr}^{-1} \cdot \text{MeV}^{-1}$ of the western coast of Europe. A Google Earth map was used.

- b For the center of the death spot, a coincidence is in effect-the direction of the geomagnetic induction vector coincides with the direction of the alpha particle intrusion-the alpha particle movement is not affected by the deflecting magnetic force. Such a coincidence occurs twice a year for latitudes in the band from 28°N to 48°N [4]. For latitudes outside this band, such a coincidence is impossible; the fast alpha particles do not reach the Earth's surface, or their flux on the surface decreases rapidly as they move away from the band.

The moment of occurrence of a flow of fast alpha particles cannot be predicted, but the dates of increased risk for a given point on the Earth's surface between 30°N - 50°N can be calculated by the latitude of the location [10]. For example, for the EUROSTAT NUTS-2 region of Switzerland (Bern) (See Figure 2), with a latitude of 47°N , the dates with maximum risk are around May 24 and July 20. On these dates, the inclination (63.6°) of the geomagnetic vector for Switzerland is close in magnitude to or coincides with the culmination of the Sun (the Earth's atmosphere is thinnest at the moment of the Sun's culmination, and there is no deflecting magnetic force for alpha particles if they intrude at this time from the Sun). The increased risk of health incidents outdoors around local noon is a further argument in favor of the healthfulness of the indoor midday break ('siesta') practiced in Mediterranean countries.

The arguments given above reveal a new, previously unsuspected cosmic cause of neoplasm mortality-solar corpuscular alpha radiation. This type of radiation may be the leading external cause of neoplasm diseases. The risk, particularly for neoplasms, is highest for inhabitants in mountainous regions. During the day, the hours around local noon are the most risky for a cosmic impact that could cause subsequent illness. The risk is also increased on certain days of the year (depending on the latitude of the observation point) when the solar culmination coincides with the angle of the geomagnetic vector (inclination). If a stream of high-energy solar alpha particles were to hit the Earth at this time, there would be no deflecting magnetic force on them, making it easier for them to reach the Earth's surface.

A part of the recorded pulses of alpha particles forms an unnatural series with fixed periods (meaning a permanent meridian, both for the registration point and the death point) and relatively constant magnitudes, which allows the hypothesis of their artificial origin [10,12, 15, 23].

References

- Nikolay Takuchev (2019) Solar corpuscular radiation and mortality from various forms of ischemic heart disease in Bulgaria for the interval 2005-2015. AIP Conference Proceedings 2075: 130005.
- Takuchev N (2020) Ischemic Heart Diseases Mortality for Bulgaria Partly Depends on Solar Corpuscular Radiation? ES Journal of Cardiology 1: 1010.
- Nikolay Takuchev (2021) How Ischemic Heart Diseases Mortality Depends on Solar Corpuscular Radiation: A Case from Bulgaria. Chapter in: Challenges in Disease and Health Research 8: 44-55.
- Nikolay Takuchev (2021) Does the Angel of Death Sometimes Use Solar Alpha Particles to Take Our Souls? Book Publisher International 1-40.
- Nikolay Takuchev (2021) Solar Corpuscular Fluxes of Alpha Particles and Mortality from Ischemic Heart Disease in Bulgaria. In: Proceedings of a scientific conference Radiation Safety in the Modern World 17-18.
- Nikolay Takuchev (2022) The Solar Alpha Particles Invading Earth and Ischemic Heart Diseases-the Worldwide Killer Number One, Connected? London Journal of Research in Science https://journalspress.com/LJRS_Volume22/The-Solar-Alpha-Particles-Invading-Earth-and-Ischemic-Heart-Diseases-the-World-Wide-Killer-Number-One-Connected.pdf.
- Nikolay Takuchev (2023) The High-Energy Positive Solar Particles Invading the Earth with Contribution to the Mortality from Ischemic Heart Diseases? SCIREA Journal of Physics 8: 54-65.
- Nikolay Takuchev (2023) Alpha particle fluxes and ischemic heart disease mortality study for Bulgaria and Los Angeles, California, USA. J Mod Appl Phys 6: 1-6.
- Nikolay Takuchev (2023) Solar Alpha Particles and Death from Ischemic Heart Disease. Journal of Current Medical Research and Opinion CMRO 06: 1615-1625.
- Takuchev N (2023) Space threat to the USA – increase in mortality from ischemic heart disease due to high-energy alpha particles entering Earth's orbit. J Mod Appl Phys 6: 1-13.
- Nikolay Takuchev (2023) Solar Alpha Particles and Death from Ischemic Heart Disease. Int J Complement Alt Med. (International Journal of Complementary & Alternative Medicine) 16: 228-232.
- Takuchev N (2023) Humanity is subject to a deadly cosmic impact – is non-human intelligent intervention also possible? J Mod Appl Phys 6: 1-11.
- Takuchev N (2023) High Energy Solar Alpha Particles are Dangerous for the People on the Earth's Surface. J. Phys. Astron 11: 368.
- Nikolay Takuchev (2023) Mortality from Diseases of the Circulatory System in Europe and the Mediterranean Connected with Space Energetic Alpha Particles. Journal of Cardiology Research Reviews & Reports 4: 1-7.
- Takuchev N (2023) Human Mortality Depends on Streams of Cosmic Alpha Particles, some of which have Signs of Artificial Origin J Space Explor 12: 305.
- Takuchev N (2024) Space Alpha Particle Fluxes towards the Earth's Surface Increase Human Mortality-Examples from the Balkans and Greece. Open J of Astro 2: 000105.
- Takuchev NP (2024) Streams of High-Energy Solar Alpha Particles - The Cause of Both Rising Global Temperature (Global Warming) and Partly Human Mortality? Phys Sci & Biophys J 8: 000267.
- Nikolay Takuchev (2024) Solar Alpha Particle Radiation Increases Human Mortality – an example from the HIV disease mortality in the European Union. International Journal of Medical Science and Clinical Invention 11: 7096-7105
- Takuchev N (2024) Solar Alpha Particle Radiation Increases Human Mortality – An Example from the Tuberculosis Mortality in the European Union. Recent Adv Clin Trials 4: 1-9.
- Takuchev N. Solar Alpha Particle Radiation Increases Human Mortality – An Example from the Ischemic Heart Diseases Mortality in the European Union. Cardiol Vasc Res. 2024; 4(2): 1-9.
- Nikolay Takuchev (2024) Solar Alpha Particle Radiation Increases Human Mortality -An Example from the Malignant Melanoma Mortality in the Europe and Mediterranean. Biomed J Sci & Tech Res <https://biomedres.us/pdfs/BJSTR.MS.ID.008901.pdf>.
- Nikolay Takuchev (2024) Solar Alpha Particle Radiation Affects Human Psyche and Behavior-Examples of Mental and Behavioral Disorders Mortality in the European Union. Biomed J Sci & Tech Res <https://biomedres.us/pdfs/BJSTR.MS.ID.008994.pdf>.
- Takuchev N (2024) Space Alpha Particle Fluxes Towards the Earth Increase Human Mortality. Is It Possible an Intervention of Non-Human Intelligence? J Med - Clin Res & Rev 8: 1-19.
- Nikolay Takuchev (2024) The Malignant Non-Hodgkin's Lymphoma and the Cutaneous Melanoma, have a Common Risk Factor? Aren't these the Solar Alpha Particle Streams? Biomed J Sci & Tech Res <https://biomedres.us/pdfs/BJSTR.MS.ID.009085.pdf>.
- Nikolay Takuchev (2024) Is it Possible that the Connection Between Solar Lentigines and Type 2 Diabetes Mellitus Can Result from a Common Cause – Solar Alpha Particle Streams? Biomed J Sci & Tech Res <https://biomedres.us/pdfs/BJSTR.MS.ID.009196.pdf>.
- EUROSTAT, European short list of deaths by cause, death rate by NUTS 2 region of residence Statistics | Eurostat (europa.eu) 10/10/2023.
- Data from GOES satellites <https://www.ncei.noaa.gov/data/goes-space-environment-monitor/access/avg/>.
- Lakin GF (1990) Biometrics. Moscow. Higher School Publishing House 208-214.
- PSTAR <https://physics.nist.gov/PhysRefData/Star/Text/PSTAR.html>.

30. ASTAR <https://physics.nist.gov/PhysRefData/Star/Text/ASTAR.html>.
31. Data on the geomagnetic field <http://www.intermagnet.org/data-donnee/dataplot-eng.php?type=xyz>.
32. Cosmic Ray Data Base (CRDB), charged cosmic rays. <https://tools.ssdc.asi.it/CosmicRays/>.

Copyright: ©2025 Nikolay Takuchev. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.