

## Comparison of Hematological Indices in Patients before and After Coronary Artery Bypass Grafting (CABG)

Hamid Hoseinikhah<sup>1</sup>, Daryoush Hamidi-Alamdari<sup>2</sup>, Parisa Asadyan-sohan<sup>3</sup>, Omid Javedanfar<sup>1</sup>, Farshad Samadie<sup>1</sup>, Hamed Banpou<sup>3</sup>, Ali Javandoost<sup>3</sup>, Aliasghar Moeinipour<sup>1</sup> and Malihe Aghasizadeh<sup>4\*</sup>

<sup>1</sup>Department of Cardiac Surgery, Imam Reza Hospital, Mashhad University of Medical Sciences, Mashhad, IR Iran

<sup>2</sup>Vascular and Endovascular Surgery Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>3</sup>Department of Metabolic Syndrome research center, Mashhad University of Medical Sciences, Mashhad, IR Iran

<sup>4</sup>Department of Internal Medicine, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

### ABSTRACT

**Subject:** Clinically, a reduction in hematocrit (Hct) or hemoglobin (Hb) levels is a notable condition commonly observed in patients undergoing coronary artery bypass grafting (CABG). Postoperative anemia is a risk factor for mortality in patients undergoing CABG, affecting both the early and late stages of recovery. However, the underlying mechanisms remain incompletely elucidated. We aimed to investigate the relation between the hematologic indices with two surgery methods (on-pump & off-pump).

**Methods:** Fifty consecutive CABG cases were randomly assigned to group I (On-pump, n= 24) and group II (Off-pump, n = 26). General clinical and surgery parameters, hematologic incidence were measured in two blood samples, the first sample was taken from each patient on the morning of the day of surgery after 12-hr fasting, and the second sample on the first postoperative day at 24-hr after surgery. Statistically assessing serum levels pre- and post-CABG and comparing between the two groups was analyzed and the correlation between parameters was evaluated.

**Result:** In the baseline, there were significant differences between the on-pump and off-pump groups regarding heart rate (HR) ( $p = 0.002$ ). Investigation after surgery showed that the difference in HR and urea levels in pre- and post-CABG was significant when compared between the two groups. A significant difference was observed in the white blood cell, it became less after the operation. The decrease in platelet and PT level in patients who underwent the on-pump and off-pump surgery (diff = -23.5 and diff = -0.62, respectively) was shown and this difference remained significant between the two groups.

**Conclusion:** We achieve the on-pump method places greater stress on blood cells, resulting in reduced levels of white blood cell in patient outcomes. It can be suggested that it should be adjusted for patients undergoing both on-pump and off-pump surgeries.

### \*Corresponding author

Malihe Aghasizadeh, Ph.D, Department of Internal Medicine, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

**Received:** August 20, 2024; **Accepted:** August 30, 2024; **Published:** September 06, 2024

### Introduction

In a significant number of developed and developing countries, coronary artery disease (CAD), largely induced by atherosclerotic in the coronary arteries, is recognized as a major cause of death (1). Managing patients suffering from CAD is challenging due to the complexity inherent in the disease (2). To treat patients suffering from these diseases (regardless of drug treatments) surgical intervention is frequently essential, and in most cases, cardiopulmonary bypass (CPB) is needed. Nonetheless, a range of post-surgical complications including heart failure, multiple organ failure (MOF), respiratory issues, as well as liver and kidney failure can occur (3-5).

Coronary artery bypass grafting (CABG) ranks among the most frequently executed surgical procedures, with around 200,000 operations conducted each year in the United States. In Western Europe, the incidence of CABG is approximately 62 per 200,000 individuals (6). Throughout the past century, the procedure has

undergone significant evolution, shifting from a high-risk operation to a relatively secure option (with the current mortality rate ranging between 1% to 2%) (7). Estimates indicate that in Iran, the number of patients requiring CABG each year surpasses ten thousand (8). Even so, the procedure is notably complex and is linked to various complications (7, 9). In this method, the final objective is to bypass a stenosed or occluded coronary artery (9). This surgical approach can be carried out with arterial or venous grafts, with or without cardiopulmonary bypass, and endoscopic harvesting of conduits (10). Conventionally, CABG is executed through the method of cardiopulmonary bypass, known as on-pump CABG. This technique involves halting the heart's function and employing a heart-lung machine to sustain adequate circulation throughout the body (11). Over the last few years, the off-pump CABG method has increasingly become a favored alternative technique in the field of cardiac surgery. Surgeons utilize an artificial circulatory system, known as the heart-lung machine, to replace the functions of the heart and lungs during operations. This strategy involves

bypass grafts on a heart that is actively beating and helps mitigate the inflammatory response typically linked with CPB (12).

Clinically, a decline in hematocrit (Hct) or hemoglobin (Hb) levels is an important condition often observed in patients undergoing CABG (13-15). Studies have demonstrated a relationship between mean platelet volume (MPV) and the occurrence of late saphenous vein graft disease in patients who have undergone CABG (16). A research investigation led by Utku Unal and colleagues demonstrated a significant correlation between preoperative MPV and hematocrit levels, as well as the occurrence of adverse outcomes following CABG. Conversely, the analysis of white blood cell counts, including differential leukocyte counts, revealed no significant correlation with negative events post-CABG (17). Existing literature suggests that an increase in preoperative red blood cell distribution width (RDW) independently forecasts mortality and the occurrence of early complications to CABG (18-21). However, the findings obtained relied on a single preoperative RDW assessment, and there is insufficient information regarding the influence of postoperative changes in RDW (22). According to Urbanowicz et al., neutrophils (among the various postoperative inflammatory indices) emerge as one of the most significant predictors of long-term survival after off-pump coronary artery bypass surgery, especially when evaluated in conjunction with preoperative characteristics (23). Despite various research, there is still limited information available about the levels of hematological parameters before and after surgery. Therefore, the upcoming study aims to investigate blood parameters in patients before and after CABG.


## Patients and Methods

### Patients

This study was conducted at the Imam Reza Hospital in Mashhad, specifically in the Department of Cardiac Surgery. Fifty patients aged between 40 and 80 years, who sought coronary artery bypass surgery and provided written consent, were consecutively enrolled. The study received ethical approval from the Mashhad University of Medical Sciences (IR.MUMS.IRH.REC.1403.065) and informed consent was obtained from all participants. The selection criteria for CABG surgery were based on guidelines from the American College of Cardiology/American Heart Association Task Force on Practice Guidelines, focusing on patients with stable angina pectoris. Two groups of patients were created, on-pump or off-pump groups according to surgical technique that was determined by the cardiac surgeon. Of 50 patients, twenty-four patients in the group underwent CABG using the on-pump technique, while twenty-six patients had the off-pump approach. Patients typically take medications such as Osivix, Plavix, and ASA. They were instructed to discontinue these medications five days prior to the operation. Additionally, some patients with high blood pressure were prescribed drugs like Metoprolol, Inderal, Losartan, Captopril, Nitroglycerin, and Nitrocantin.

### Inclusion and Exclusion of Criteria

Patients (aged between 40 and 80 years), who refer to the Imam Reza Hospital in Mashhad and undergoes CABG were consecutively enrolled. The following criteria excluded patients from participating in the study: 1) those without written consent, 2) patients with an EF < 20%, 3) individuals with preoperative kidney disease indicated by a serum creatinine > 0.3 mg/dL, 4) patients with coagulation disorders prior to surgery, 5) individuals with concurrent valve and coronary diseases, 6) The need to perform unanticipated complex surgery during the operation, 8) Death of the patient in the operating room, Flow diagram of the study of the study was shown in Figure 1.



**Figure 1:** Flow diagram of the Study.

### Measurements

Upon patient enrollment, essential demographic information is collected: age (in years) and sex. A unique case number is assigned for the surgery and the operation date is recorded. The type of vascular disease is categorized as either 1VD (single-vessel disease), 2VD (two-vessel disease), 3VD (three-vessel disease), or LM (left main coronary artery disease). Additionally, the Ejection Fraction (EF, %) is measured to assess left ventricular function. Pre-existing cardiovascular risk factors are then evaluated. 1) Hypertension is defined as systolic blood pressure (SBP)  $\geq 140$  mmHg, diastolic blood pressure (DBP)  $\geq 90$  mmHg, or the current use of antihypertensive medications. 2) diabetes mellitus (DM), based on a fasting blood sugar level of  $\geq 126$  mg/dL, or if the patient is currently taking diabetic medications. 3) Chronic conditions like chronic renal failure (CRF), 4) chronic obstructive pulmonary disease (COPD) is screened for, with their presence or absence recorded. 5) dyslipidemia is assessed based on total cholesterol  $\geq 200$  mg/dL, LDL-C  $\geq 130$  mg/dL, or the current use of cholesterol-lowering medications.

### Intraoperative and Postoperative Monitoring

Upon ICU admission, meticulous monitoring commences to identify and address potential postoperative complications. Blood loss is recorded at specific time points: 1, 6-, 12-, 24-, and 48-hours following surgery. The occurrence of atrial fibrillation (AF) and ventricular arrhythmias (VT/VF), is tracked. Additionally, any reoperations or blood transfusions required are documented.

The Vasopressor-Inotropic Score (VIS) is calculated as follows: Vasopressor-Inotropic score =  $([\text{Dopamine} + \text{Dobutamine}] \times 1) + (\text{Milrinone} \times 15) + ([\text{Epinephrine} + \text{Norepinephrine} + \text{Isoproterenol}] \times 100)$ , and is recorded upon ICU admission and at regular intervals throughout their stay (6-, 12-, 24-, and 48-hours post-admission) to assess their hemodynamic stability. Hemodynamic parameters (systolic, diastolic, and mean arterial pressures (MAP) along with heart rate (HR) are continuously monitored and recorded at the same intervals as the VIS.

Arterial blood gas measurements are taken pre-CPB, every 30 minutes during CPB, immediately upon ICU admission, and again at 12- and 24-hours post-admission. CPB parameters are documented, including aortic cross-clamp time and total CPB duration. Additionally, inotropic support needs within the first 2 hours after surgery and the use of an intra-aortic balloon pump (IABP) are recorded. Mortality rates are tracked during hospitalization and up to one month after discharge. Surgical site infections (SSIs), categorized as superficial or deep (with potential dehiscence), are monitored. The duration of mechanical ventilation and length of stay in both the ICU and the hospital are

documented. We note any difficulties encountered during extubating (weaning process) within 48 hours post-surgery. Neurological complications, if any, are recorded. Blood samples are drawn to measure serum glucose and liver enzyme levels preoperatively, upon ICU admission, and 24 hours later.

### Statistical Analysis

The patients' information was collected, sorted and processed and then entered into SPSS version 20 software. The description of the data was expressed with the help of descriptive statistics indicators in the form of frequency and mean + standard deviation and in the form of suitable tables and graphs. After checking the normality of the data, T-test or Mann-Whitney-U was used to check the relationship between the qualitative and quantitative variables, and the relationship between the qualitative variables was performed using the chi-square test. Also, in order to control intervening variables, logistic regression model was used to determine the relationship between the independent variable and the categorized

values of the dependent variables. A statistically significant level of 5% was considered.

### Result

#### Baseline Characteristics of Patients

The findings from the initial analysis revealed the baseline characteristics of the patients presented in Table 1. Among the 50 patients, there were 15 females (30%) and 35 males (70%). The average age was  $59.92 \pm 10.71$  years. The subjects were divided into two groups: On-pump (N=24) and Off-pump (N=26). The distribution of men and women was balanced in both groups. Significant differences were observed between the on-pump and off-pump groups regarding heart rate, with means of  $90.63 \pm 20.64$  and  $74.42 \pm 14.06$ , respectively ( $p = 0.002$ ). However, no notable differences were found in other variables between the two groups. There were not significant differences between the on-pump and off-pump groups in terms of cross clamp and CPB and postoperative mechanical ventilation duration.

**Table 1: Characteristics of Patients Undergoing Coronary Artery Bypass Surgery (CABG) and Comparing it in Two Methods, on Pump and Off Pump.**

Variable	On pump, N=24	Off pump, N=26	P-value
Characteristics			
Age	58.67±11.93	61.08±9.6	0.4
Female	7(29.2)	8(30.8)	0.9
Male	17(70.8)	18(69.2)	
CVD Type			
1VD	-		0.6
2VD	5(20.8)	7(26.9)	
3VD	14(58.3)	16(61.5)	
4VD	5(20.8)	3(11.5)	
Discription of Operation			
ICU Duration	2.96±2.17	2.50±0.99	0.35
SBP	111.38±15.03	111.31±17.62	0.98
DBP	64.83±12.23	62.42±10.67	0.46
Heart Rate	90.63±20.64	74.42±14.06	0.002
EF %	46.2±9.5	49.3±11.4	0.3
Cross Clamp Duration (min)	49.27±17.83	-	-
CPB Duration (min)	73.04±20.84	-	-
Duration of Mechanical Ventilation (hours)	9.42±3.3	9.65±5.4	0.8

### Comparing pre- and Post-CABG

The evaluation of hematological incidence before and after coronary artery bypass grafting (CABG) are displayed in Table 2. This table presents the baseline parameter levels and their differences (labeled as "diff") 24 hours after surgery, along with the statistically significant variations observed. As indicated, there was a significant difference in MCHC levels before and after the operation in patients who underwent on-pump surgery. In this group, a significant decrease was observed in platelet and PT ( $-23.5 \pm 41.14$  and  $-0.62 \pm 1.15$ , respectively). However, patients who underwent off-pump surgery showed an increase in PT.

As can be seen from the result of white blood cell indices, a significant difference was observed in the serum levels of WBCs and lymphocytes after surgery in both groups. What is interesting in this data is that decrease difference in lymphocytes level in on-pump group was two-fold more than off-pump ( $-8.2 \pm 15$ ,  $-4.8 \pm 5.58$ , respectively).

Data are shown as Mean ± SD for Normal Variables and Median (Interquartile Range ) for Non-Normal Variables. Abbreviations:

**Table 2: Comparing the Hematology Markers in Patients before and after CABG and Comparing it in two Methods, on Pump and Off Pump.**

Variable	On pump (N=24)			Off pump (N=26)			
	Baseline	Diff	P-value	Baseline	Diff	P-value	*P value
White Blood Cell Indices							
WBCs (103/M)	15.15±6.4	-2.55±4.3	0.008	13.84±4.5	-2.01±4.2	0.025	0.001
Lymphocytes	14.02±14.9	-8.2±15	0.015	13.9±4.3	-4.8±5.58	0.001	0.94
Neutrophils	81.22±2.63	7.8±1.2	0.06	89.25±3.6	12.3±3.2	0.2	0.154
INR	1.19±1.21	-0.2±0.19	0.615	1.6±2.25	-0.49±2.2	0.280	0.814
RED Blood Cell Indices							
RBC (103/M)	3.86±0.71	-0.23±0.62	0.085	4.02±0.46	-0.009±0.55	0.931	0.153
Hgb (g/dL)	10.97±1.7	-0.49±1.76	0.1	11.5±1.8	-0.34±3.3	0.903	0.394
Hct (%)	10.97±4.73	-2.02±4.8	0.054	34.2±4.6	-0.107±4.7	0.908	0.005
MCV (fL)	85.02±5.5	-0.62±1.5	0.062	84.8±4.3	-0.046±2.003	0.907	0.001
MCH (pg)	28.59±2.1	0.28±0.9	0.147	28.6±2.1	0.39±1.17	0.099	0.001
MCHC (g/dL)	28.59±2.1	0.6±1.1	0.018	33.6±1.4	0.51±1.5	0.106	0.011
RDW (%)	10.87±3.53	-2.02±0.2	0.153	11.45±2.53	-1.02±2.5	0.210	0.59
Platelet indices							
Platelet (103/uL)	194.2±53.7	-23.5±41.14	0.01	241.3±75.9	-13.42±44.47	0.136	0.001
MPV (fL)	10.19±1.21	-0.2±0.19	0.615	9.6±0.25	-0.49±2.2	0.280	0.410
PDW (fL)	45.39±7.56	1.2±0.18	0.525	49.5±2.25	2.48±3.8	0.580	0.851
PT	14.9±1.4	-0.62±1.15	0.015	13.83±1.5	4.3±25.01	0.389	0.002
PTT	38.75±4.3	-4.65±4.03	0.220	38.50±6.1	-4.6±5.1	0.168	0.562

\* P value of difference between diff in two group. The P-value less than 0.05 was considered statistically significant. Data are shown as Mean ± SD for normal variables and Median (Interquartile Range ) for non-normal variables.

**Abbreviations:**

**WBC:** White Blood Cell; **MID:** Mid-Range Absolute Count; **RBC:** Red Blood Cell; **HGB:** Hemoglobin; **HCT:** Hematocrit; **MCV:** Mean Cell Volume; **MCH:** Mean Cell Hemoglobin; **MCHC:** Mean Cell Hemoglobin Concentration ; **RDW:** Red Blood Cell Distribution Width; **MPV:** Mean Platelet Volume; **PDW:** Platelet Distribution Width.

T Student and Mann whitney Tests were used for Analysis. P-value < 0.05 is Considered Statistically Significant.

**Comparing on-Pump and Off-Pump CABG**

Table 2 illustrated the comparison of parameters levels between the on-pump and off-pump methods, before and after CABG. The last column in the table indicates the p-value for the variance between “diff” in the on-pump and off-pump groups.

Comparison of the changes of WBC as post- minus pre-operation data between on- and off-pump CABG surgery was demonstrated that there were decreasing changes in group 1 and group 2. These differences were significant in both groups and also P value of difference between diff in WBC in studied groups was remained significant (P=0.001).

In addition, P value of difference between diff in MCV, MCH, MCHC, and Hct in studied groups was remained significant. As can be seen from the table, A significant decrease was observed in the levels of Platelet (-23.5±41.14) and PT (-0.62±1.15) in on pump group. Also, P value of difference between diff in these parameters in studied groups was remained significant.

**Discussion**

A systematic evaluation of peripheral blood hematologic parameters following cardiac surgery is essential for monitoring changes and implementing necessary corrections. Anemia is a

common condition characterized by a deficiency of red blood cells or hemoglobin in the blood, leading to decreased oxygen transport to tissues (24). Its prevalence varies based on the population and underlying health conditions. In the context of patients undergoing CABG (coronary artery bypass grafting), studies have shown that anemia can be observed in over 90% of cases, highlighting its significant impact on this patient group (25). This can be attributed to acute blood loss occurring during and after surgery. Approximately 85% of intraoperative and early postoperative bleeding is related to technical factors and is caused by CPB, which leads to hemodilution.

As mentioned in the literature review, postoperative anemia is a risk factor for mortality in patients undergoing CABG, both in the early and late stages (26). According to the primary objective of this project, we have monitored changes in hematologic incidence throughout the operation to gain a deeper understanding of anemia before and after the CABG intervention. Blood loss is a frequent issue in cardiac surgery, often necessitating additional interventions, particularly in the case of significant blood loss (27). One of the important issues that emerges from our findings that was not investigated in other studies, is decreasing the level of WBC after 24 hours operation in patients who underwent surgery with on-pump method is more reduced than in patients who were operated

with off-pump method significantly. A possible explanation for this might be that high WBC count prior to CABG surgery serves as an independent risk factor for ischemic events occurring one year following the surgery (28). Despotis et al. (22) indicated that patients exhibiting the highest increase in white blood cell (WBC) count during cardiopulmonary bypass (CPB) are most prone to bleeding in the first 24 hours post-surgery. On-site WBC count measurements can act as an indicator of the inflammatory response to CPB, aiding in the identification of patients who are at an elevated risk for significant bleeding. With this insight, physicians may be able to apply anti-inflammatory medications and blood conservation strategies to manage the situation (29).

Our study was demonstrating a significant difference in lymphocytes and neutrophil in both group of patients who underwent the on-pump and off-pump surgery. It is encouraging to compare our investigation with that found by point out the hematological changes in patients undergoing coronary artery bypass surgery. They showed the average levels of leukocytes and neutrophils rose quickly, reaching their peak on day 2, whereas the average lymphocyte count dropped sharply, reaching its lowest point on day 1 after surgery (+74.7%, +127.1%, and -52.4% respectively compared to preoperative values,  $p < 0.001$ ) (30).

Prior studies have not noted the importance of difference between kind of CABG surgery. We found that a significant decrease was observed in the levels of Platelet and PT in on pump group and these parameters in studied groups was remained significant. Lako (2015) showed that the average platelet count decreased to the lowest value on day 2 after surgery from the preoperative (30). The acute decrease in hematocrit (Hct) led to a reversible impairment of platelet function, which was evident by an increase in bleeding time (BT) and a reduction in the Thromboxane B2 level in shed blood at the template BT site (31). In most patients, platelet counts typically rise thereafter, returning to pre-surgery levels around postoperative days 5 to 7 (32). Miyauchi et al. observed a significant decrease in platelet count upon the start of bypass, with this low level persisting until the third postoperative day (33).

## Conclusion

Our findings showed that decreasing the level of WBC after 24 hours operation in patients who underwent surgery with on-pump method is more reduced than in patients who were operated with off-pump method significantly. The average values of leukocytes in pre and post operation increased, while this difference was not significant between on- and off- pump. We found that a significant decrease was observed in the levels of Platelet and PT in on pump group and these parameters in studied groups was remained significant.

## Declaration Section

### Informed Consent

Informed consent was obtained from all individual participants included in the study.

### Authors' Contributions

All authors were involved in the interpretation of the data and contributed to writing the manuscript. All authors read and approved the final manuscript.

### Availability of Data And Materials

The data that support the findings of this study are available from the corresponding author upon reasonable request.

### Ethics Approval

The approval of the ethics committee of the Mashhad University of

Medical Sciences (IR.MUMS.IRH.REC.1403.065) was received and informed consent form.

### Consent for publication

Not applicable.

### Acknowledgements

We are grateful for the cooperation of the participants of MASHAD Study, without whom this research would not have been conducted.

### Financial Support

None.

### Conflict of Interest

The authors have no conflict of interest to disclose

### References

1. Malakar AK, Choudhury D, Halder B, Paul P, Uddin A, et al. (2019) A review on coronary artery disease, its risk factors, and therapeutics 234: 16812-16823.
2. Dourado LOC, Bittencourt MS, Pereira AC, Poppi NT, Dallan LAO, et al. (2018) Coronary artery bypass surgery in diffuse advanced coronary artery disease: 1-year clinical and angiographic results 66: 477-482.
3. Pacholewicz J, Kuligowska E, Szylińska A, Walerowicz P, Biskupski A, et al. (2023) The rate of postoperative mortality and renal and respiratory complications are increased in patients with extreme obesity undergoing cardiac surgery—a Retrospective Observational Cohort Study of 8848 Patients 1155-1166.
4. Tanner TG, Colvin MOJL (2020) Pulmonary complications of cardiac surgery 198: 889-896.
5. Cizmic A, Rahmanian PB, Gassa A, Kuhn E, Mader N, et al. (2023) Prognostic value of ascites in patients with liver cirrhosis undergoing cardiac surgery 18: 302.
6. Melly L, Torregrossa G, Lee T, Jansens J-L, Puskas JDJJotd (2018) Fifty years of coronary artery bypass grafting 10: 1960.
7. Selim MJNEjom (2007) Perioperative stroke 356: 706-713.
8. Ebrahimi M, Sabouri Arteh A, Rahimi HRJRiCM (2015) Coronary arteries bypass grafting stenosis 2: 32-36.
9. McKhann GM, Goldsborough MA, Borowicz LM, Mellits ED, Brookmeyer R, et al. (1997) Predictors of stroke risk in coronary artery bypass patients 63: 516-521.
10. Hussain SMA, Harky AJJjomr (2019) Complications of coronary artery bypass grafting 6: 1-5.
11. Kirk KC, Aldridge RA, Sistino JJ, Zellner JL, Crumbley AJ, et al. (2001) Coronary artery bypass grafting with and without cardiopulmonary bypass: a comparison analysis 33: 86-90.
12. Larmann J, Theilmeier GJBp, anaesthesiology rC (204) Inflammatory response to cardiac surgery: cardiopulmonary bypass versus non-cardiopulmonary bypass surgery 18: 425-438.
13. Zindrou D, Taylor KM, Bagger JPJTL (2002) Preoperative haemoglobin concentration and mortality rate after coronary artery bypass surgery 359: 1747-1748.
14. Fang WC, Helm RE, Krieger KH, Rosengart TK, DuBois WJ, et al. (1997) Impact of minimum hematocrit during cardiopulmonary bypass on mortality in patients undergoing coronary artery surgery 96: II-194.
15. Ferraris VA, Ferraris SPJTJot, surgery c (1996) Risk factors for postoperative morbidity 111: 731-741.
16. Tavil Y, Şen N, Yazici HU, Hizal F, Açıkgöz SK, et al. (2010) Relationship between elevated platelet volume and saphenous vein graft disease 33: 161-167.
17. Unal EU, Ozen A, Kocabeyoglu S, Durukan AB, Tak S, et

- al. (2013) Mean platelet volume may predict early clinical outcome after coronary artery bypass grafting 8: 1-7.
18. Taşoğlu I, Turak O, Özcan F, Nazlı Y, Kadiroğullari E, et al. (2013) Preoperative red cell distribution width as a predictor of mortality in patients after coronary artery bypass grafting 16: 1-6.
  19. Ünal EU, Özen A, Kubat E, Sarıtaş AJJCS (2013) Red cell distribution width is associated with early adverse events following coronary bypass surgery 1: 43-46.
  20. Benedetto U, Angeloni E, Melina G, Pisano C, Lechiancole A, et al. (2013) Red blood cell distribution width predicts mortality after coronary artery bypass grafting 165: 369-371.
  21. Warwick R, Mediratta N, Shaw M, McShane J, Pullan M, et al. (2013) Red cell distribution width and coronary artery bypass surgery 43: 1165-1169.
  22. Lee SI, Lee SY, Choi CH, Park C-H, Park KY, et al. (2018) Relation between changes in red blood cell distribution width after coronary artery bypass grafting and early postoperative morbidity 10: 4244.
  23. Urbanowicz T, Michalak M, Olasińska-Wisniewska A, Rodzki M, Witkowska A, et al. (2022) Neutrophil counts, neutrophil-to-lymphocyte ratio, and systemic inflammatory response index (SIRI) predict mortality after off-pump coronary artery bypass surgery 11: 1124.
  24. Beris P, Munoz M, García-Erce JA, Thomas D, Maniatis A, et al. (2008) Perioperative anaemia management: consensus statement on the role of intravenous iron. *British journal of anaesthesia* 100: 599-604.
  25. Kim P, Dixon S, Eisenbrey AB, O'Malley B, Boura J, et al. (2007) Impact of acute blood loss anemia and red blood cell transfusion on mortality after percutaneous coronary intervention. *Clinical cardiology* 30: 35-43.
  26. Van Straten AH, Soliman Hamad MA, Van Zundert AJ, Martens EJ, Schönberger JP, et al. (2009) Preoperative hemoglobin level as a predictor of survival after coronary artery bypass grafting: a comparison with the matched general population. *Circulation* 120: 118-125.
  27. Karkouti K, Wijeyesundera DN, Yau TM, Beattie WS, Abdelnaem E, et al. (2004) The independent association of massive blood loss with mortality in cardiac surgery. *Transfusion* 44: 1453-1462.
  28. Rashidi F, Jamshidi P, Kheiri M, Ashrafizadeh S, Ashrafizadeh A, et al. (2012) Is leukocytosis a predictor for recurrence of ischemic events after coronary artery bypass graft surgery? A cohort study. *International Scholarly Research Notices* 2012: 824730.
  29. Despotis GJ, Levine V, Goodnough LT (1997) Relationship between leukocyte count and patient risk for excessive blood loss after cardiac surgery. *Critical care medicine* 25: 1338-1346.
  30. Lako S, Dedej T, Nurka T, Ostreni V, Demiraj A, et al. (2015) Hematological changes in patients undergoing coronary artery bypass surgery: a prospective study. *Medical Archives* 69: 181.
  31. Valeri CR, Cassidy G, Pivacek LE, Ragno G, Lieberthal W, et al. (2001) Anemia-induced increase in the bleeding time: implications for treatment of nonsurgical blood loss. *Transfusion* 41: 977-983.
  32. Warkentin TE, Levine MN, Hirsh J, Horsewood P, Roberts RS, et al. (1995) Heparin-induced thrombocytopenia in patients treated with low-molecular-weight heparin or unfractionated heparin. *New England Journal of Medicine* 332: 1330-1336.
  33. Miyauchi Y, Isomura K (1982) Quantitative and functional changes in platelets and their preservation during extracorporeal circulation. *The Journal of Cardiovascular Surgery* 23: 383-390.

**Copyright:** ©2024 Malihe Aghasizadeh, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.