

Research Article

Open Access

Diagnosis of the Syndrome of Excessive Bacterial Growth of the Small Intestine in Patients with Bauhinia Insufficiency and After its Surgical Correction (First Results)

Vladimir Leonidovich Martynov* and Natalia Vladimirovna Kazarina

Headquarters of the State Educational Establishment of the City of Nizhny Novgorod, Russia

ABSTRACT

Objective: To determine the syndrome of excessive bacterial growth in the small intestine (SIBO) in patients with insufficiency of the bauhinia valve (NBZ) and after its surgical correction.

Methods of examination: Patients of the studied groups were examined using direct and indirect methods of diagnosis of SIBO. Bacteriological examination of ileal aspirate and other surgical material was performed. All patients underwent a hydrogen respiratory test with a load of lactulose and performed a qualitative reaction of urine to indican.

Characteristics of the material: 50 patients were examined, of which 30 were included in the main group, who underwent surgical correction of NBZ - Bauginoplasty; 20 patients are included in the control group in which the ileocecal valve is consistent. Patients of the main group were examined before surgery and on the 7th and 45th day after Bauginoplasty.

Main results: All patients of the main group had SIBS of varying severity, in 80% of patients SIBS was localized in the distal small intestine. Patients with normal function of ileocecal valve syndrome of excessive bacterial growth did not suffer. Intraoperatively, 76% of patients showed signs of mesenteric mesentery of the small intestine, and the fact of bacterial translocation in SIBO was also confirmed. After 7 days after surgical correction of the Bauhinia valve, the normalization of peak and background excretion of hydrogen was observed in 37% of patients. For 45 days in all patients the hydrogen curve corresponded to the norm.

Conclusions: the deficiency of the bauhinia valve is obligately accompanied by the syndrome of excessive bacterial growth in the small intestine, surgical correction is an effective method for correcting the syndrome of excessive bacterial growth in patients with NBD.

*Corresponding author

Martynov VL, Headquarters of the State, Educational Establishment of the City of Nizhny Novgorod, Russia, E-Mail: hirurgia12@mail.ru

Received: September 26, 2020; **Accepted:** October 03, 2020; **Published:** October 30, 2020

Keywords: Excess Bacterial Growth Syndrome, Bauhinia Valve Insufficiency, Hydrogen Respiratory Test, Bauhinoplasty

Introduction

The deficiency of the Bauhinium Damper (NBZ) is widespread among the population. In the study of 900 protocols of irrigoscopy, performed for various indications and not related to the diagnostic search for a Bauhinia flap insufficiency, the NBZ was detected in 64% of patients as an accompanying finding [1].

Widespread NBZ among the population of Russia due to connective tissue dysplasia (DST) and an extensive range of diseases of the colon and terminal part of the small intestine, determining, respectively, the primary and secondary NBZ [2].

Diseases of the gastrointestinal tract (GIT) occupy a prominent place in the structure of the pathology of internal organs, in Russia they are about 2500 per 100 thousand population [3]. In recent years, much attention has been paid to functional diseases of the

digestive organs, which is primarily due to their high prevalence and the costs incurred by health care in connection with the examination and treatment of such patients [4]. Leading experts say that irritable bowel syndrome (IBS) affects from 10 to 20% of the adult population of developed countries, which is up to 50% of all visits to a gastroenterologist [5]. According to recent studies, the frequency of DST among adolescents reaches up to 20% (in the cohort about 3000 studied, according to the data of Professor Nechaeva G. I., and 8.5% in a sample of 400 adults [6].

According to our data, classical appendectomy, and even more so appendectomy with technical difficulties and appendectomy under conditions of a complicated course of acute appendicitis, can lead to an insolvency of the bauhinia valve [1,2]. The fact of intimate anatomical and functional proximity of the valve structures of the ileocecal junction and appendix is undeniable. Appendectomy destroys these connections, and leads to the development of secondary acquired NBZ.

Many authors admit the fact of ascending retrograde infection of the small intestinal mucosa at NBD [7-9]. A pathological condition in which the titer of microorganisms in the mucous membrane of the small intestine exceeds 105 CFU / ml, or colonization of the colonic flora occurs, has been called Small Bacterial Overgrowth Syndrome (SIBO).

SIBR is a pathological condition that develops as a result of bacterial contamination of the small intestine by various microflora and is accompanied by functional disorders of the digestive conveyor [10, 11].

Direct evidence of the occurrence of SIBR in case of the failure of the Bauhinia Damper (NBZ) is the experience of Larry S Miller (2012) [9]. In a clinical experiment, the author modeled the ileocecal valve insufficiency by placing a 4-lumen probe behind the lips of the bauhinia valve in the direction of the ileum of 19 healthy volunteers during a colonoscopy. 1 month after the manipulation with the help of the hydrogen breathing test, all the subjects under study revealed a syndrome of excessive bacterial growth of the small intestine. Thus, by virtually destroying the Bauhinia flap, the author achieved the emergence of SIBR from volunteers.

Currently, SIBO is recognized as a key pathogenetic mechanism in the development and persistence of many diseases of the digestive tract and associated extra-digestive conditions (bronchial asthma, dermatological atopy, diabetes, autoimmune allergic conditions, arthritis, obligate precancerous conditions, etc.).

The actual prevalence of SIBO is unknown today. Clinically, this pathological condition is poorly diagnosable due to the non-specificity of its symptoms, which are often attributed to the main disease that causes SIBO [12]. In fact, SIBO is extremely common in the presence of one or more predisposing pathological conditions. Thus, Ardatskaya MD (2011) indicates that the frequency of detection of excessive bacterial growth in the small intestine in various diseases (chronic gastritis, peptic ulcer, chronic cholecystitis, pathology of the hepatobiliary system, inflammatory and other intestinal diseases (in tons irritable bowel syndrome), scleroderma, neuropathy (as a complication of diabetes mellitus), the effects of surgery, etc.) is 40–99% [8]. The results of the last meta-analysis, which included 11 studies, showed that pathological respiratory tests are much more frequently recorded in patients with IBS than in healthy individuals (OR = 4.46; 95% CI = 1.69 - 11 , 80) [13]. Similar results were achieved in the second meta-analysis, published a year later, - pathological test results were significantly more frequent in patients with IBS: odds ratio (OR) 4.46, 95% confidence interval (CI) 1.69–11.80. At the same time, taking into account age and gender showed an even more significant result (OSH 9.64, 95% CI 4.26–21.82).

According to the literature, the problem of SIBR is currently a purely therapeutic one. Treatment of bacterial overgrowth syndrome in the small intestine provides for the treatment of the underlying disease, replacement therapy for malabsorption syndrome and antibacterial therapy [10,14]. The main group of drugs used for SIBR is intraintestinal antibiotics [15-17]. It should be recognized that in the absence of an impact on the cause, successful pathogenetic treatment is temporary, which is confirmed by the example of rifaximin therapy. The frequency of recurrence of SIBR after effective therapy with rifaximin reached 44% after 9 months [18].

Many authors separately stop at an obvious conclusion - the prediction of the SIBR and the risk of its recurrence after successful antibiotic therapy depends primarily on the therapeutic effect on the primary background disease that has contributed to the contamination of the small intestine [10,19]. Gabrielli M. (2013) rightly points out that the recurrence of SIBO in the event of persisting predisposing causes often leads to a chronic process and a complex, and sometimes impossible, conservative correction [18].

Thus, we have not encountered literary data that would describe etiopathic therapy. In our opinion, the problem of SIBO in patients with functional pathology of the gastrointestinal tract in many cases is due to damage to a specific anatomical structure, namely, the Bauhinia valve, which is a surgical problem.

We have not met domestic studies on the etiopathic effect on the syndrome of excessive bacterial growth in patients with proven deficiency of the bauhinia valve. The few foreign works are devoted only to the statement of the fact that there is a SIBR at the NBZ [7,9].

The purpose of the study is to determine the syndrome of excessive bacterial growth of the small intestine in patients with proven insufficiency of the bauhinia valve and after its surgical correction.

Materials and Methods

The study was conducted on the basis of the surgical department of the city hospital "Nizhny Novgorod City Clinical Hospital No. 12" in the period from September 2014 to December 2014. The main group included 20 patients (16 women, 8 men, age from 23 to 57 years) with proven by the results of irrigoscopy the insufficiency of the bauhinia valve. All patients of the main group performed Bauginoplasty according to the methods of prof. Martynova V.L. A control group of 20 patients was also formed (11 women, 9 men, age from 28 to 49 years), who underwent an irrigoscopy in order to diagnose diseases of the large intestine, according to the results of which the ileo-cecal valve turned out to be healthy. Simultaneously, patients of the main group underwent appendectomy. The removed vermiform process was sent for histological examination. In the case of intraoperative diagnosis of mesenteric mesentery of the ileum, the altered lymph node was resected, a backseed from a section of the lymph node was performed, and his histological examination was performed.

Preparation for irrigoscopy was carried out for two days and consisted in the prohibition of food intake containing slags, in the formulation of high enemas with a total volume of 1.5 to 3 liters 3-4 times a day. It is imperative to thoroughly empty the cecum from the intestinal contents, since its presence in the form of an embossed mass may prevent the radiopaque substance from passing into the ileum with the failure of the bauhinia valve. This can lead to a false negative conclusion. The taut intestine must be filled with a radiopaque substance. Repeated X-ray examination after a bowel movement is fundamentally important, since during bowel movement a maximum pressure is created in the intestine, which, if the ileocecal locking apparatus fails, will be accompanied by a pronounced reflux of the radiopaque substance into the ileum (Fig. 1.).




Figure 1: Throwing barium into the ileum with irrigoscopy

We performed intraoperative digital diagnostics of the failure of the bauhinia valve. Normally, the gap of the thin-colic transition is no more than 2 cm and does not miss the first finger of the surgeon's left hand. In the NBZ environment, on average, this transition has a diameter of 3–3.5–4 cm, which determines the almost unhindered passage of 1 finger of the left hand (so conveniently) by the surgeon into the ileum. The technique consists in the following: after extracting the ileocecal angle from the right ileal fossa, the surgeon 1 finger of the left hand brings the lateral wall of the ascending colon, opposite the ileocecal anastomosis, closer to this anastomosis and tries to insert it and the finger into the ileum. With its introduction into the ileum, in our opinion, the diagnosis of NBZ is established (Fig. 2.). If it is impossible to perform this manipulation, the diagnosis of the NBZ can be rejected.



Figure 2: The first finger of the left hand of the surgeon with the lateral wall of the ascending colon is inserted into the ileum. Intraoperative digital diagnosis NBZ installed

Direct intraoperative diagnostic method for small intestine SIBR is a bacteriological examination of aspirate of the intraluminal and mucosal contents of the distal ileum (Fig. 3.). The technique of material sampling is as follows: 20 cm proximal to the ileocecal junction after isolation of the 10-cm long loop of the ileum with a length of 10 cm by ileum intestinal tissue, the lumen puncture is performed. The needle is held obliquely relative to the intestinal wall, 10 ml of sterile physiological NaCl solution is injected into the intestinal cavity. The exposure is maintained for 5 minutes, the contents of the isolated ileum are aspirated, until the needle passes through the mucous layer of the intestine, the needle is removed. The puncture site is peritonized. Aspirate is sent for bacteriological examination.



Figure 3: Intraoperative sampling of the intraluminal and mucosal contents of the distal ileum followed by bacteriological examination

Indirect techniques included the determination of urinary indican (by the method of Obermeyer) before the operation and on the 7th and 45th day after bauginoplasty; hydrogen respiratory test (VDT) with a lactulose load, which was carried out using a Gastrolyser apparatus before the operation and on the 7th and 45th day after the operation. A preoperative study of patients with NBZ using these techniques was performed prior to the radiopaque examination of the intestine, or 4 weeks after irrigoscopy. Patients of the main group did not receive antibiotic therapy in the postoperative period. Preoperative preparation consisted of 3-fold cleansing enemas, which could not affect the biocenosis of the small intestine.

In order to determine the syndrome of excessive bacterial growth in the patients of the control group, indirect diagnostic methods were carried out, which included the determination of urine indican (by the method of Obermeier) and VDT with a load of lactulose, performed using a Gastrolyser apparatus. The study preceded the irrigoscopy.

Preparation of patients for VDT with lactulose was carried out according to the method proposed by [20]. The diagnostic criterion for the presence of SIBS in patients was an increase in H₂ excretion after exercise in the period from 0 to 60 min by 10 ppm (in the absence of abdominal symptoms after lactulose, such as abdominal pain, rumbling, flatulence, diarrhea) and 5 ppm (in the presence of abdominal symptoms). The group included patients with basal secretion of hydrogen up to 5 ppm. It was carried out using statistics.

Results and Discussion

Postoperative complications in the patients of the main group were not observed. The proposed preoperative irrigoscopy. It was confirmed that a chronic inflammatory catarrhal was observed in all 20 cases.

Intraoperative mesadenitis was detected in 16 patients of the main group. It has been shown that there has been no clear link between the cortical and cerebral layers. The following is the result: *E. coli* hemolytic was determined in 9 cases (104 CFU / ml), *St. aureus* - in 7 cases (103-4 CFU / ml).

It has been taken for 20 patients who have been diagnosed in the biocenosis. The spectrum of microorganisms is determined by the following microorganisms: *E. coli* hemolytic was determined in 10 patients (105-107 CFU / ml), *St. aureus* - in 6 patients (106-108 CFU / ml), *C. albicans* - in 4 patients (105-109 CFU / ml). The sown flora is identical.

Bauginoplasty (BP) according to the results of the patient's patient's life cycle (Fig. 3), the colony peak at 90th minute).




Figure 3: Results of excretion of H₂ in patients of the main group before surgery

The patients of the main group were distributed as follows (Table 1).

Table 1: The distribution of patients in the main group (n = 20)

Criteria	Number of patients
The degree of SIBR of the small intestine (Mechetina TA, 2011)[26]	
degree: increase by 20-50 ppm	4
degree: increase by 50-100 ppm	14
degree: increase by 100 ppm and more	2
Localization of SIBR (Gabrielli M., 2013) [18].	
proximal SIBR	2
distal SIBR	14
total SIBR	4

The determination of urinary indican in patients of the main group before surgical correction gave positive results in 16 cases (Table 2). The presence of an insignificant (1 degree - increase up to 50 ppm) distal syndrome of excessive bacterial growth in 4 patients in whom urine indican was negative may indicate a lack of sensitivity of this screening test compared to VDT.

Table 2: The results of the determination of indican urine with varying degrees of SIBR

The degree of SIBR of the small intestine (Mechetina TA, 2011) [26]	Urine indican (according to Obermeier)
1 degree: increase by 20-50 ppm	“Negative” (less than 100 mg / ml in the blood)
2 degree: increase by 50-100 ppm	“Positive” (more than 100 mg / ml in the blood)
3 degree: increase by 100 ppm	and more “Positive” (more than 100 mg / ml in the blood)

On the 7th and 45th days of the postoperative period, in the absence of antibiotic therapy, indican urine in all operated patients (n = 20) was negative.

According to the results of VDT with lactulose load in patients of the main group on the 7th day after the Bauginoplasty (BP) (Fig. 4), all of the patients underwent a significant reduction in the amount of hydrogen released, while the lactulose passage

through the intestines normalized (the large intestinal peak is determined at 75 minute). On the 45th day after Bauginoplasty, VDT corresponded to the norm in all patients (Fig. 5).




Figure 4: Results of H₂ excretion in patients of the main group on the 7th day after surgery.




Figure 5: Results of H₂ excretion in patients of the main group on the 45th day after surgery

Fig. 5. Results of H₂ excretion in patients of the main group on the 45th day after the operation. The urine index in the control group of patients with the proven consistency of the bauhinia valve was confirmed with an irrigoscopy. The results of the hydrogen respiratory test in the control group (Fig. 6) did not differ from the normal values described in the literature [20].




Figure 6: Results of H₂ excretion in control group patients

Our studies confirm the presence of small intestine SIBR in patients with NBZ, which is consistent with the data of foreign

authors and fits into the logic of colonic reflux pathogenesis in the small intestine [7,9]. On the reliability of the data in this study can not speak. However, an obvious difference is found in the results of VDT in patients of the compared groups: a significant decrease in the concentration of hydrogen released after the operational benefit (Fig. 7.). The first facts obtained objectively indicate the effectiveness of Bauginoplasty, the positive clinical results of which we observed in 500 operated patients with NBZ [21].




Figure 7: Comparative characteristics of H₂ excretion in patients of the control and main groups before and after bauginoplasty

The results of the bacteriological examination of the ileum aspirate of patients with NBZ confirm the presence of SIBR detected by HTD with a load of lactulose. At the same time, a correspondence was established between the degree of SIBR (according to the results of VDT) and the degree of contamination of ileal aspirate (Table 3). This is consistent with the data of E. Mendoza (2007), which established the sensitivity (85.7%) and specificity (90.9%) of the hydrogen breath test with lactulose compared with the “gold standard” [22].

Table 3: The ratio of the degree of SIBO and the degree of dissemination of ileal aspirate in patients with NBZ (n = 20)

The degree of SIBR (Mechetina TA, 2011) (ppm)	Seediness of ileal aspirate (CFU / ml)	Number of patients
1 degree	to 10^6	4
2 degree	10^7	14
3 degree	to 10^{8-9}	2

The determination of indican urine makes it possible to suspect SIBS in patients with a pronounced degree of dissemination of the small intestine (increased H₂ excretion above 50 ppm), but not informatively in patients with SIBS grade 1 (elevation of H₂ to 50 ppm) compared to VDT. The revealed diagnostic value of this method retains its value as a screening test for small intestine SIBR of 2 and 3 degrees.

The phenomena of mesadenitis and the presence of microorganisms in the lymph nodes confirm the translocation of microorganisms through the intestinal wall. The fact of bacterial translocation diagnosed in patients with SIBO of the small intestine under NBZ conditions is consistent with the literature data [23-27].

References

- Martynov VL (2013) Surgical treatment of reflux disease: Monograph / V.L. Martynov. - Nizhny Novgorod, Publishing house LLC Center for operational printing 535.
- Martynov VL (2006) Refluxes of the digestive tract and their surgical correction: dis. ... Dr. honey. Sciences: 14.00.27 / Martinov Vladimir Leonidovich. - Saransk 261.
- Filippova TV (2014) The value of medical genetic counseling for diseases of the digestive system / T. V. Filippova // RZhGK 1: 56-61.
- Sheptulin AA (2010) Gastroesophageal reflux disease and functional diseases of the gastrointestinal tract: is there any connection? / A. A. Sheptulin, M. A. Wiese-Hripunova // RZhGGK. - 2010. - T.20 4: 44-48.
- Sperber AD, Grossman DA, Quigley EM (2012) The Global Foundation for World Gastroenterology Organization symposium // Am. J Gastroenterol 107: 1602-1609.
- Nechaeva GI, Yakovlev VM, Gromova OA, Torshin IYu, et al. (2009) Connective tissue dysplasia in children and adolescents. Innovative hospital-saving technologies for diagnosis and treatment in pediatrics. M.: Union of Pediatricians of Russia 96.
- Oren Zaidel (2003) Uninvited Guests: The Henry C. Lin. // Practical Gastroenterology 27: 24-37.
- Ardatskaya MD (2011) Syndrome of excess bacterial growth: a tutorial / M. D. Ardatskaya. - Moscow: Forte print 56.
- Larry S Miller (2012) Ileocecal valve dysfunction in small intestinal bacterial overgrowth: A pilot study / Larry S. Miller, Anil K. Vege, Aiswarya Madanam Sampath, Shital Prabhu, Sesha Krishna Kotapati, Kian Makipour // WGJ 18: 6801-6808.
- Curly Yu A (2014) Interrelation of irritable bowel syndromes and excessive bacterial growth: is it there? / Yu. A. Kucheryavy, S.V. Cheryomushkin, E.A. Mayevskaya, E.A. Sutugina // RZGGK 2: 5-14.
- Maev IV (2011) Diagnosis and treatment of the syndrome of excessive bacterial growth in the small intestine / I. V. Maev, N. Yu. Ivashkina, Yu. A. Kucheryavy, TS Oganesyan // Eksper. wedge. gastroenterol 3: 125-129.
- Bures J (2010) Small intestinal bacterial overgrowth syndrome / Bures J., Cyrany J., Kohoutova D., Forst L.M., Rejchr T.S., Kventin A.J., Vorisek V., Kopasova M. // World J Gastroenterol 28: 2978-2990.
- Ford AC (2009) Small intestinal bacterial overgrowth in irritable bowel syndrome: review and meta-analysis / Ford A. C., Spiegel B. M., Talley N. J., Moayyedi P. // Clin Gastroenterol Hepatol 7: 1279-1286.
- Avdeev V (2010) Diagnosis and treatment of bacterial overgrowth syndrome in the small intestine / V. Avdeev // Doctor 12: 1-3.
- Plotnikova E Yu (2013) Some aspects of diagnosis and treatment of excessive bacterial contamination of the small intestine in clinical practice [electronic resource] / E. Yu. Plotnikova, M. V. Borsch, M. V. Krasnova, E. N. Baranova // Attending doctor 4.
- Poluektova EA (2013) Treatment of irritable bowel syndrome from the standpoint of modern ideas about the pathogenesis of the disease / E. A. Poluektova, S. Yu. Kuchumova, A. A. Sheptulin, V. T. Ivashkin // RZGGK 1: 57-65.
- Kucheryavyi Yu A (2010) Syndrome of excess bacterial growth / Yu. A. Kucheryavyi, T.S. Oganesyan // RGGK 5: 63-68.
- Gabrielli M (2011) Prevalence of small intestinal bacterial overgrowth in Parkinson's disease / Gabrielli M., Bonazzi P., Scarpellini E., Bendi AE, Lauritano EC, Fasano A., Ceravolo MG, Capecci M., Rita Bentivoglio A., Provinciali L., Tonali PA, Gasbarrini A. // Mov Disord 26: 889-892.
- Lauritano E (2008) Small intestinal bacterial overgrowth recurrence after antibiotic therapy / Lauritano E., Gabrielli M., Scarpellini E., Lupascu A., Novi M., Sottillii S., Vital EG, Cesarino V., Serricchio M., Cammarota G., Gasbarrini G., Gasbarrini A. // Am J Gastroenterol 103:

2031-2035.

20. Ledochowski M (2008) Hydrogen Breath tests. 1st Edition / M. Ledochowski. — Innsbruck: «Akademie» 59.
21. Gabrielli M (2013) Diagnosis of small intestinal bacterial overgrowth in the clinical practice / M. Gabrielli, G. D. Angelo, T. D. I. Rienzo, E. Scarpellini, V. Ojetto // European Review for Medical and Pharmacological Sciences 17: 30-35.
22. Mendoza E (2007) Diagnosis of small intestinal bacterial overgrowth in children: C., Crismatt C., Matos R. // Biomedica 27: 325-332.
23. Wiest R (2005) Bacterial translocation in cirrhosis / Wiest R., Garcia-Tsao G. // Hepatology 41: 422-433.
24. Hashimoto N (2002) Effect of portal portal hypertension on gut mucosa / Hashimoto N., Ohyanagi H. // Hepatogastroenterology 49: 1567-1570.
25. Heuman DM (2004) Persistent ascites and low sodium, identify patients with MELD scores and D. Abu-assi, S. G., Habib A. // Hepatology 40: 802-810.
26. Zharkova MS (2012) Influence of excess bacterial growth syndrome and bacterial translocation on the course of the disease in patients with cirrhosis of the liver / M.S. Zharkova, M.V. Mayevskaya, V.T. Ivashkin // RGGG 5: 56-63.
27. Mechetina TA (2011) Syndrome of excessive bacterial growth in the small intestine after cholecystectomy: dis. ... Cand. honey. Sciences: 14.01.28 / Tatyana Anatolevna Mechetina. - Moscow 134.

Copyright: ©2020 Martynov VL. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.