

Case Report

Open Access

Invasive Papillary Carcinoma of the Breast in a Male: Case Report and Review of the Literature

Mohamed Reda El Ochi^{1*}, Massine El Hammoui², Mohamed Amine Essaoudi¹, El Hassane Kabiri² and Mohamed Oukabli¹

¹Department of Pathology, Mohamed V Military Hospital, Mohammed V University, Hay Riad, Rabat, Morocco

²Department of thoracic surgery, Mohamed V Military Hospital, Mohammed V University, Hay Riad, Rabat, Morocco

Abstract

Breast tumors are commonly found in women. Hereby we report a case of invasive papillary carcinoma of the breast in a 65 years old male who presented at our department with 6 months history of left breast pain. Ultrasonographic (US) examination showed a well demarcated nodule measuring 1,8 cm of diameter. A biopsy was performed and the pathological examination revealed an invasive carcinoma not otherwise specified. After discussion the patient underwent modified radical mastectomy, which showed invasive solid papillary carcinoma. The patient has been well on adjuvant chemotherapy without any recurrence for 6 months.

*Corresponding author

Mohamed Reda El Ochi, Department of Pathology, Mohamed V Military Hospital, Mohammed V University, Hay Riad, Rabat, Morocco.
E-mail: elochi20@yahoo.fr

Received: April 26, 2022; **Accepted:** April 30, 2022; **Published:** May 11, 2022

Introduction

Male breast tumors are extremely rare with less than 1% of all breast cancer cases [1]. Papillary carcinoma of the male represents less than 1% of all breast tumors [2]. The average age at diagnosis is 67 years [3]. The clinical presentation is frequently a painless localized mass in the subareolar region. Other clinical findings include nipple discharge or ulceration, besides axillary lymph node swelling.

As far as we know, a few cases of invasive solid papillary carcinoma of the male breast have been reported.

The prognosis of this entity is controverted.

Case Presentation

A 65 years old man with history of 45 years of smoking, diabete and hypertension presented at our department with 6 months history of left breast pain. On clinical examination, a firm and sharply demarcated nodule was found in the subareolar region of the left breast. A slightly retracted nipple was noted without ulceration or discharge. There were palpable axillary lymphadenopathies. The patient did not report any family history of breast tumor. Mammogram showed a nodule in the left subareolar region. US analysis 18x13x10 mm solid mass in the same localization. Percutaneous biopsy was performed and an invasive mammary carcinoma was reported. The patient underwent modified radical mastectomy with a complete dissection of axillary lymph nodes (Figure.1).

Macroscopically, the excised tumor was 2 x 1,5 x 1 cm in size. The nipple was slightly retracted without skin ulceration (Figure.1). Cut surface showed well defined solid cystic mass. 11 lymph nodes were removed.

Microscopically, the tumor showed an invasive solid papillary carcinoma reported as grade II according to the modified Bloom Richardson without lymph node involvement (Figure. 2&3).

Immunohistochemically, myoepithelial cells were not seen on the invasive component (Figure 4). Tumor cells were strongly positive for estrogen and progesterone receptors.




Figure 1: (A) Preoperative position of the Patient. (B) Postoperative view, MP:Major Pectoralis Muscle, CN: Charle-Bel Nerve, AV: Axillary Vein, ND: Nerve of Major Dorsalis. (C) Gross Examination of the Mastectomy Specimen




Figure 2: Expansile Nodules Composed of Solid Epithelial Proliferation Punctuated by Delicate Fibrovascular Cores (Hemalum& Eosin x 100)




Figure 3: Invasive Component with Destructive Growth (Hemalun&Eosin x 400)




Figure 4: P63 Antibody Showing the Absence of Myoepithelial Cells in the Invasive Component (Immunohistochemistry x 250)

As a result of the above findings, a final diagnosis of invasive solid papillary carcinoma was made.

Discussion

The male mammary glands have a discoid shape with a diameter similar to that of the areola, 3 to 4 mm thick, they are composed of fatty tissue with some ducts and connective tissue, but without the development of acini and lobules. This breast tissue can respond to hormonal stimuli, which results in the growth of connective tissue and conduits (gynecomastia), and can also develop cancer [4].

Male breast carcinomas are rarely observed and it accounts for 0,6 – 1 % of all mammary cancers [5].

The cause of male breast cancer is poorly understood, but hormonal imbalance, obesity and some testicular abnormalities are implicated in this disease [6]. Other risk factors include advanced age, radiation exposure, smoking and chronic heat exposure, which may disturb testicular function [7]. A positive family history and BRCA2 gene mutation confer a high risk of male breast cancer [8]. In our case only smoking and advanced age are found as risk factors.

It has been documented that male breast cancer is somewhat more likely to be diagnosed in left breast than right. Most tumors were confined to the central subareolar area.

Solid papillary carcinoma is commonly seen on mammography as oval or round and well demarcated lesion, exceptionally it can show obscure margin [9].

The predominant histologic subtype is invasive carcinoma not otherwise specified. Solid papillary carcinoma is extremely rare. Histologically, the tumor shows solid sheets, jigsaw pattern with more ragged and irregular margins, coupled with absence of myoepithelial cells is considered invasive carcinoma, thin fibrovascular cores are interspersed within the tumor, the cells are plasmacytoid, uniforms with speckled chromatin.

In majority of cases, strong estrogen and progesterone receptor expression was demonstrable and half of cases showed neuroendocrine differentiation [10].

Regarding the prognosis, solid papillary carcinoma without invasive component is not associated with metastatic axillary involvement [11]. When invasive component is present, some metastatic lesions have been reported in the literature [12].

The treatment of choice consists on partial or radical mastectomy, adjuvant chemotherapy can be added according to the staging information [13]. Radiation and endocrine therapy are still under investigation [14].

Conclusion

To conclude, solid papillary carcinoma of the breast is an uncommon disease in women. Here, we have described a rare case of this entity with an invasive component in a male patient which must raise the concern when dealing with a painless mass in male breast. Much work in term of molecular characterization of male breast cancer is needed to provide advice for optimal treatment choices.

Competing Interests

The authors declare that they have no conflicts of interest.

References

1. Cutuli B, Le-Nir CC, Serin D, Kirova Y, Gaci Z, et al. (2010) Male breast cancer. Evolution of treatment and prognostic factors. Analysis of 489 cases. Crit Rev Oncol Hematol 73: 246-254.
2. Wang-Rodriguez J, Cross K, Gallagher S, Djahanban M, Armstrong JM, Wiedner N, et al. (2002) Male breast carcinoma: correlation of ER, PR, Ki-67, Her2-Neu, and p53 with treatment and survival, a study of 65 cases. Mod Pathol 15: 853-881.
3. Mathew J, Perkins GH, Stephens T, Middleton LP, Yang WT (2008) Primary breast cancer in men: clinical, imaging, and pathologic findings in 57 patients. AJR Am J Roentgenol 191: 1631-1639.
4. Hittmair AP, Lininger RA, Tavassoli FA (1998) Ductal carcinoma in situ (DCIS) in the male breast. A morphologic study of 84 cases of pure DCIS and 30 cases of DCIS associated with invasive carcinoma - a preliminary report. Cancer 83: 2139-2149.
5. Giordano SH, Cohen DS, Buzdar AU, Perkins G, Hortobagyi GN (2004) Breast carcinoma in men: a population-based study. Cancer 101: 51-57.
6. Misra SP, Misra V, Dwivedi M (1996) Cancer of the breast in a male cirrhotic: is there an association between the two? Am J Gastroenterol 91: 380-382.
7. Mabuchi K, Bross DS, Kessler II (1985) Risk factors in male breast cancer. J Natl Cancer Inst 74: 371-375.
8. Thorlacius S, Tryggvadottir L, Olafsdottir GH, Jonasson JG, Ogmundsdottir HM, et al. (1995) Linkage to BRCA2 region in hereditary male breast cancer. Lancet 346: 544-545.
9. Schneider JA (1989) Invasive papillary breast carcinoma: mammographic and sonographic appearance. Radiology 171: 377-379.
10. Pal SK, Lau SK, Kruper L, Nwoye U, Garberoglio C, et al. (2010) Papillary carcinoma of the breast: an overview. Breast Cancer Res Treat 22: 637-645.
11. Nassar H, Qureshi H, Adsoy NV, Visscher D (2006) Clinicopathologic analysis of solid papillary carcinoma of the breast and associated invasive carcinomas. Am J Surg Pathol 30: 501-507.
12. Saremian J, Rosa M (2012) Solid papillary carcinoma of the breast: a pathologically and clinically distinct breast tumor. Arch Pathol Lab Med 136: 1308-1311.
13. Fentiman IS, Fourquet A, Hortobagyi GN (2006) Male breast cancer. Lancet 367: 595-604.
14. Y. Otsuki, M. Yamada, S. Shimizu, Kaori Suwa, Masayuki Yoshida, et al. (2007) Solid-papillary carcinoma of the breast: clinicopathological study of 20 cases Pathol Int 57: 421-429.

Copyright: ©2022 Mohamed Reda El Ochi, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.