

Short Communication

Open Access

Remodeling of Nasal Mucosa in Patients with Rhinitis

Nelson Augusto Rosario Filho*, Loreni C Siqueira Kovalhuk and Ederaldo Queiroz Telles

Federal University of Paraná, Curitiba, Brazil

*Corresponding author

Nelson Augusto Rosario Filho, Rua Padre Camargo 453, Curitiba, Brazil.

Received: September 26, 2023; **Accepted:** September 28, 2023; **Published:** September 30, 2023

Rhinitis is a chronic inflammatory process of the nasal mucosa that can progress to structural alterations characterized as airway remodeling such as observed in asthmatic patients [1]. Remodeling is the result of recovering from an inflammatory damage and represents the relationship between an initial inflammation and the subsequent tissue damage. Although not well defined in rhinitis it could be related to increased thickness of basement membrane, pseudofibrosis and epithelial detachment in histologic samples of nasal mucosa [2].

The nasal mucosa is a physical barrier to particles, allergens and pollutants and a conditioner for inspired air. The nasal pseudostratified columnar epithelium is separated from the lamina propria by a continuous basement membrane. Chronic inflammation could induce structural changes in the epithelium or lamina propria [3].

Samples of nasal mucosa can be easily obtained allowing the study of pathological alterations of the allergic reaction [4]. Nasal lavage is a relatively noninvasive and easy to perform technique for the quantitative measurement of cell distribution, inflammatory mediators and effect of treatment [5,6]. The simultaneous analysis of the cell infiltrate and the degree of cell activation of nasal secretion and nasal mucosa biopsy samples show differences between the two compartments [7].

Our aim was to study the inflammatory process in samples simultaneously obtained by nasal lavage (NL) and from the nasal mucosa of atopic and nonatopic patients with rhinitis.

Fifty-six young adults (median age of 24.5 years) and a diagnosis of allergic rhinitis (n = 36) and non-allergic rhinitis (n = 20) were submitted to allergy skin tests to common inhalant allergens, nasal lavage (NL) with measurement of albumin and interleukin-8 levels, total and differential cell counting by microscopy. Mucosal histopathology determined the extent of epithelial lesion, and degree of basement membrane thickening. All patients with rhinitis had nasal obstruction had undergone correction of deviated nasal septum and (or) turbinate hypertrophy and were examined before the surgical procedures. Nasal lavage technique was previously described [6].

Nasal mucosa samples from the inferior turbinates were obtained at partial turbinectomy or by mucosal biopsy of the anteroinferior tip of the turbinate and analysed by light microscopy. The degree of

epithelial lesion (EL) and basement membrane thickening (BMT) were based on a score proposed by Ponikau et al [8].

Symptom-adjusted score (pruritus, sneezing, nasal obstruction, nasal secretion/sniffing, post-nasal secretion/snorting) and sign-adjusted score (mucosa color, turbinate volume increase, aspect and volume of nasal secretion), were graded on a scale from 0 to 3 (absent, mild, moderate and severe), with a maximum score of 24 [9]. The total symptom score was higher in atopic subjects (9 [1-18]) than in nonatopics (6.5 [0-12]) (p = 0.01). Total and differential cell counts, as well as albumin and IL-8 levels, are shown in Table 1. The percentage of eosinophils was higher among atopic patients (p < 0.01).

Table 1: Total and differential cell count, albumin and IL-8 levels in nasal lavage of atopic and non-atopic subjects

Nasal lavage	Atopic (n=36)	Non-Atopic (n=20)	P value
Total cells	127,000 (10×10^3 - 6.134×10^3)	128,000 (24×10^3 - 682×10^3)	0,90
Neutrophils (%)	41.5 (0-87)	17.5 (0-83)	0,24
Eosinophils(%)	3 (0-66)	1 (0-21)	<0.01
Epithelial cells (%)	48 (8-98)	76 (10-100)	0,07
IL-8 (pg/mL)	80 (30-1300)	81.5 (30-604)	0,45
Albumin (µg/mL)	16 (5-338)	16.5 (5-105)	0,67

Median (limits); Mann-Whitney test.

The highest sensitivity and specificity for eosinophil counts were 4% in the NL, a cutoff point that differentiated atopic from non-atopic subjects.

In the presence of some degree of epithelial lesion the findings of nasal lavage were similar in both groups of allergic and non-allergic subjects. There were no differences in BMT regarding the number of eosinophils in the NL. On the other hand, BMT was associated with an increase of neutrophils in the nonatopic (NAT) subjects' NL (p < 0.01). The frequency of BMT was similar in atopic (67%) and non-atopic (55%) subjects as well as the proportion of eosinophils and neutrophils in NL of both subgroups.

Therefore these findings may suggest the participation of eosinophils in basement membrane thickening of patients with allergic or non-allergic rhinitis. Activated eosinophils cause epithelial damage and loss of epithelial integrity in rhinitis patients [10]. This study showed that there were some differences in the inflammatory process, but they may result in similar structural damage to the nasal mucosa of rhinitis patients independently of the allergic status.

References

1. Bourdin A, Neveu D, Vachier I, Paganin F, Godard P, et.al. (2007) Specificity of basement membrane thickening in severe asthma. *J Allergy Clin Immunol* 119: 1367-1374.
2. Watelet JB, Van Zele T, Gjomarkaj M, Canonica GW, Dahlen SE, et.al. (2006) Tissue remodelling in upper airways: where is the link with lower airway remodeling?. *Allergy* 61: 1249-1258.
3. Watelet JB, Bachert C, Gevaert P, Van Cauwenberge P (2002) Wound healing of the nasal and paranasal mucosa: a review. *Am J Rhinol* 16: 77-84.
4. Howarth PH, Persson CG, Meltzer EO, Jacobson MR, Durham SR, et.al. (2005) Objective monitoring of nasal airway inflammation in rhinitis. *J Allergy Clin Immunol* 115: 414-441.
5. Bascom R, Pipkorn U, Lichtenstein LM, Naclerio RM (1988) The influx of inflammatory cells into nasal washings during the late response to antigen challenge. Effect of systemic steroid retreatment. *Am Rev Respir* 138: 406-412.
6. Kovalhuk LCS, Rosário NA, Carvalho A (2001) Inflammatory mediators, cell counts in nasal lavage and computed tomography of the paranasal sinuses in atopic children. *J Pediatr (Rio J)* 77: 271-278.
7. Lim MC, Taylor RM, Naclerio RM (1995) The histology of allergic rhinitis and its comparison to cellular changes in nasal lavage. *Am J Respir Crit Care Med* 151: 136-144.
8. Ponikau JU, Sherris DA, Kephart GM, Kern EB, Gaffey TA, et.al. (2003) Features of airway remodeling and eosinophilic inflammation in chronic rhinosinusitis: is the histopathology similar to asthma?. *J Allergy Clin Immunol* 112: 877-882.
9. Meltzer EO, Jalowayski AA, Vogt K, Iezzoni D, Harris AG, et.al. (2006) Effect of desloratadine therapy on symptom scores and measures of nasal patency in seasonal allergic rhinitis: results of a single-center, placebo-controlled trial. *Ann Allergy Asthma Immunol* 96: 363-368.
10. Amin K, Rinne J, Haahtela T (2001) Inflammatory cell and epithelial characteristics of perennial allergic and nonallergic rhinitis with a symptom history of 1 to 3 years' duration. *J Allergy Clin Immunol* 107: 249-257.

Copyright: ©2023 Nelson Augusto Rosario Filho, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.