

Research Article

Open Access

Sensory Evaluation and Fatty Acid Composition of Broiler Chickens Fed Diets Containing with *Prosopis Africana* Essential Oil

Alagbe* and Olujimi John

Department of Animal Science, University of Abuja, Nigeria

ABSTRACT

Exploring the use of phytochemical components in plants is one of the ways to ensure safety in food and sustainable livestock production. This experiment was designed to investigate the sensory evaluation and fatty acid composition of broiler chickens fed diets containing *Prosopis africana* essential oil. A total of 540 – one- day- old broiler chicks (Arbo Acres) were randomly distributed into 6 treatments, each containing six replicates of 15 birds. Experimental diets were adequate in all nutrients (NRC, 1994). Birds in treatment one was fed basal diet without *Prosopis africana* oil, basal diet with antibiotic growth promoter (oxytetracycline at 2.5 g/kilogram) was fed to treatment two while treatment three, four, five and six were fed basal diet supplemented with 200 mg, 400 mg, 600 mg and 800 mg per kilogram respectively. Completely randomized design was used and the experiment lasted for 56 days. Feed and fresh water were offered ad libitum and other management practices were rigorously followed. Result revealed that the composition of total monosaturated fatty acid and total polyunsaturated fatty acid were significantly ($P<0.05$) influenced by the treatments. Total saturated fatty acid (TSFA) levels decrease as the level of *Prosopis africana* oil increases across the treatments ($P<0.05$). TSFA values were highest in treatment 1, intermediate in treatment 2 and lowest in treatment three through six. Omega 3: omega 6 fatty acid and atherogenicity index were significantly ($P<0.05$) different among the treatments. *Prosopis africana* oil also influenced color, flavor, tenderness, juiciness, texture, and general acceptability of broilers meat. It was concluded that the dietary supplementation of *Prosopis africana* oil has a number of phyto-constituents that can modify the fatty acid of meat, enhancing its quality, shelf life and sensory attributes. Without having a negative impact on the health of birds. It can be added to the diet of broiler chickens up to 800 mg per kilogram.

*Corresponding author

Alagbe, Department of Animal Science, University of Abuja, Nigeria.

Received: June 19, 2023; **Accepted:** June 27, 2023; **Published:** July 03, 2023

Keywords: Phytochemicals, *Prosopis Africana* Oil, Fatty Acid, Sensory Qualities, Oxytetracycline

Introduction

The need for the animal sector to create ever-more-sophisticated strategies for producing animal protein in increasingly environmentally and economically responsible ways is growing. In addition to setting up ethical and environmentally friendly structures for subsequent generations, this is done to ensure that food is safe for consumption today [1, 2]. The incorporation of feed additives to accomplish these objectives has grown increasingly crucial as regulations and consumer interest demand that nutrition for animals be gain more accurate than ever, particularly in light of the worldwide prohibition on the use of antibiotic growth promoters that was implemented in 2009 by the European Union and other nations [3, 4]. Phytochemical compounds are the most viable class of feed additives among antibiotic replacements.

Due to their wide spectrum of efficiency and additional benefits on sustainability and safety, phytochemical feed additives are expected to have a bright future in animal nutrition [5]. Phytochemical feed additions include a variety of plants, including herbs, spices, and essential oils obtained from plants, including *Prosopis africana* oil [6-8]. Because of the variety of their constituents and other impacting variables like location, harvest, species, stage, and

storage conditions, *Prosopis africana* essential oils exhibit some diversity in their chemical composition. Numerous constituents of essential oils have antibacterial properties [9-11]. For instance, Prosogerin A and B in *Prosopis* essential oil work alongside the cell membrane to create their antibacterial effects. The linking of an ion through the cell membrane results from this contact, which alters the shape of the membrane structure [12-14].

Observed that essential oil might modify the fatty acid composition of the flesh of broiler chickens and relieve the growth depression caused by a challenge with *Clostridium perfringens*, in accordance with findings [15-20]. *Prosopis africana* oil supplementation up to 800 mg/kg was found to enhance growth performance, boost the number of faecal lactobacilli, lower *Escherichia coli* counts, and stimulate animal immunological responses, according to Numerous studies suggest beneficial impacts on the morphology of the digestive tract in chickens. Birds fed *Prosopis africana* oil supplemented diets showed increased trans-epithelial resistance of the duodenal mucosa [21-28].

Prosopis africana essential oil has a wealth of potentials and contains naturally occurring bioactive substances. Presuming that it cannot be hazardous to animals at specific amount is incorrect. Modulating the meat's polyunsaturated fatty acid will extend its shelf life and contribute to a decrease in the

prevalence of cardiovascular ailments which can be achieved by the supplementation of essential oil in the diet of birds [9, 10]. Therefore, this study was undertaken to evaluate the effects of dietary supplementation of Prosopis africana essential oil on sensory evaluation meat quality of broiler chickens.

Materials and Methods

Experimental Site

The Department of Animal Science, Faculty of Agriculture, University of Abuja Teaching and Research Farm, Main Campus, located along Airport Road, Gwagwalada, Abuja, Nigeria was where this study was conducted. The Gwagwalada Area Council's administrative center is in Gwagwalada, which is situated between latitudes 8o571 and 8o551N and longitudes 7o051 and 7o061E (NPC, 2006).

Gathering, Identifying a Sample of Plants and Prosopis Africana Oil Extraction

Prosopis africana seeds used in this investigation were from the Gwagwalada area. The seeds were recognized and confirmed by the Department of Crop Protection Herbarium, University of Abuja, Gwagwalada, Nigeria, with voucher code ANS - 2021D.

The steam-distillation process, which also requires the use of an electronic scale, a round-bottom flask, water from distillation, a heating mantle composed of glass fibre, a measuring cylinder, and a separatory funnel, was used to extract Prosopis africana oil. A round-bottom flask was filled with 250 mL of distilled water and 50 g of pulverized Prosopis seed. The condenser was set up above the flask with a circular bottom after the mixture was transferred to a glass yarn heating mantle and heated to a temperature of eighty degrees centigrade. The mixture is forced to boil for a period of fifteen minutes, after which the distillate is gathered in a beaker until no more oil drips are visible, and then it is poured into a separatory funnel to yield Prosopis africana oil.

Diets and Other Aspects of Animal Care

540 mixed-sex broiler chicks (Arbo Acres) were randomly distributed into 6 treatments, each containing six replicates of 15 birds. The birds were bought from a well-known commercial hatchery in Oyo State, Nigeria. In a galvanized battery cage with semi-open pens measuring (100 cm × 75 cm × 50 cm) (length depth height), galvanized hand feeders and automatic nipple drinkers were installed. When the chicks arrived, they were measured to determine their average beginning body weight before being fed anti-stress (Vitalyte wsp + glucose) at a rate of 2 grams of glucose to 5 liters of water. A totally random experimental design was used in this study. Diets were created based on the nutritional requirements for broilers published by NRC (1994), which are displayed in Table 1. The trial diet contains 23.30% crude protein (CP), 21.40% CP, and ME at the starter phase (1-28 days) and finisher phase (29-56 days) of 2991.5 Kcal/kg and 3108.1 Kcal/kg, respectively. All birds had unlimited access to food and water for the eight-week period of the trial. Vaccination schedule is presented in Table 2.

The experimental set –up is shown as follows:

First-line treatment (T1) Prosopis africana oil -free basic diet

2nd-line treatment (T2): oxytetracycline 2.5 g/kg added to the standard diet

3rd - line treatment: basal diet plus 200 mg per kg Prosopis africana oil

Fourth-line treatment (T4): basal diet plus 400 mg per kg Prosopis africana oil

Fifth-line treatment (T5): basal diet plus 600 mg per Prosopis africana oil

Sixth-line treatment (T6): basal diet plus 800 mg/kg Prosopis africana oil

Data Collected

Meat Sample Fatty Acid Content

Twelve birds per group had their breast and thighs removed, and the overall lipids from all these parts was obtained using Bligh and Dyer's method (1959). According to Christie's description, 0.01 % sulfuric acid solution in dry methanol was used for 14 hours to produce the methyl esters of the total lipids that were extracted by sample preparation thin layer chromatography (1973). Gas-liquid chromatography examination employing a chromatograph CSI 200 fitted with a capillary column (DM-2330:30 m 0.25 mm 0.33 m) with hydrogen as a carrier gas was used to assess the fatty acid contents of lipid content. The temperature of the detector and injectors was 230 oC, and the oven temperature was first set to 160 oC for 0.2 min. After that, it was increased to 220 oC at a speed of five oC min⁻¹ and maintained there for 5 min comparing the retention periods of the benchmarks to the methyl esters allowed for their identification. By contrasting the comparable chain length with that of genuine fatty acid methyl esters, the peaks were identified. With the use of the Agilent gas chromatography chemstation software, peak areas were calculated automatically (Gujarat Technologies, USA). The fatty acid concentrations are expressed as a proportion of the overall amount of peaks that were recorded for each sample and were recognized.

Sensory Assessment

10 panelists conducted a sensory assessment of cooked samples of minced broiler chicken breast and thigh meat from six birds per treatment (from birds used for fatty acid investigation). The panelists assessed a number of factors, including color, juiciness, flavor, tenderness, and overall acceptability. Every representative of the panel received a tagged sample of beef, which was distributed one by one. After consuming every meat sample, each participant rinsed their mouth completely with warm water to prevent a baggage effect. Using only a nine-point hedonic scale, the panelists assigned scores: (i) Very dislike; (ii) Extremely dislike (iii) Medium dislike (iv), Minor dislike (v), Middle dislike (vi), Minor dislike (vii), Moderate dislike (viii), Extreme dislike (ix), and like slightly.

Proximate Analysis of Experimental Feed

Analysis of feed of experimental feed was done using Phoenix 5000 near infra-red (NIR) feed analyzer with the following specifications; wavelength (1100 – 2500 nm), photometric noise (< 15 µAu in standard range), measurement mode (reflectance or transmittance), wavelength accuracy (< 0.3 nm for Blue Sun selected published peak positions of NIST SRM 1920a standard), wavelength reproducibility (0.02 nm based on Blue Sun Scientific specific diagnostic test), analysis time (10 to 60 seconds), ambient humidity (< 85 % RH) and detector (high performance InGaAs extended ranged detector).

Statistical Investigation

Using SPSS (25.0), one-way analysis of variance was performed on all data, and Duncan multiple range tests were used to identify significant means (Duncan, 1955). If P< 0.05, significance was deemed to exist.

Table 1: Ingredient Composition of The Experimental Diets

Components	Starters mash (0-4 weeks)	Finishers mash (5-8 weeks)
Maize	52.00	60.00
Wheat offal	2.50	5.00
Soya bean meal	30.00	25.00
Groundnut cake	8.00	4.00
Fish meal (72%)	2.00	2.00
Limestone	1.50	1.50
Bone meal	3.00	3.00
Lysine	0.20	0.20
Methionine	0.20	0.20
*Premix	0.25	0.25
Salt	0.30	0.30
Toxin binder	0.10	0.10
Total	100.0	100.0
Determined analysis (% DM)		
Crude protein	23.30	21.40
Crude fibre	4.18	5.01
Ether extract	4.03	4.47
Calcium	1.50	1.60
Phosphorus	0.58	0.66
Energy (Kcal/kg)	2900.3	

*Premix supplied per kg diet: - vit A, 13,000 I.U; vit E, 5mg; vit D3, 3000I.U, vit K, 3mg; vit B2, 5.5mg; Niacin, 25mg; vit B12, 16mg; choline chloride, 120mg; Mn, 5.2mg; Zn, 25mg; Cu, 2.6g; folic acid, 2mg; Fe, 5g; pantothenic acid, 10mg; biotin, 30.5g; antioxidant, 56mg (starter's mash)

**Premix supplied per kg diet: - vit A, 9,000 I.U; vit E, 10mg; vit D3, 1500I.U, vit K, 3.8mg; vit B2, 10 mg; Niacin, 15mg; vit B12, 10mg; choline chloride, 250mg; Mn, 5.0mg; Zn, 56mg; Cu, 1.6g; folic acid, 2.8mg; Fe, 5.1g; pantothenic acid, 10mg; biotin, 30.5g; antioxidant, 56mg (finisher's mash)

Table 2: Vaccination Programme for Birds

Age / day	Vaccines	Route of administration
7	Lasota (1 st dose)	Drinking water
11	Gumboro (1 st dose)	Drinking water
15	Lasota (2 nd dose)	Drinking water
21	Gumboro (2 nd dose)	Drinking water
28	Gumboro (3 rd dose)	Drinking water

Results

The Impact of Prosopis Africana Oil on The Chicken Breast Flesh Table 3 shows the impact of Prosopis africana oil on the chicken broiler's breast muscle. Myristoleic acid, palmitic acid, stearic acid, arachidic acid, behenic acid, oleic acid, elaidic acid, erucic acid, vaccenic acid, linolenic acid, -linolenic acid, eicosapentaenoic acid, docosahexaenoic acid, and linoeladic acid values varied from 12.7 to 22.1 percent, 5.02to 11.1 percent, 1.73 – 4.10 percent, 0.10 – 0.30 percent, 1.25 – 3.94 percent, 2.01 – 4.03 percent, 10.9 – 23.7 percent, 1.00 – 1.93 percent, 0.10 – 1.02 percent, 11.7 – 25.1 percent, 0.61 – 1.34 percent, 2.80 – 13.9 percent, 1.17 – 3.21 percent, 0.42 – 1.64 percent and 0.10 – 1.72 percentcorrespondingly. Omega 6: Omega 3 (2.60–3.11 percent), total saturated fatty acids [TSFA; 16.94–53.80 percent], unsaturated fatty acids [USFA; 36.20–83.06 percent], with artheriogenic index (0.25–0.92 percent). Whereas USFA readings appeared statistically ($P<0.05$) different between treatments, TSFA values were considerably ($P<0.05$) higher in T1 and T2, midrange in T3, with smallest in T4, T5, as well as T6.

Table 3: The Impact of Prosopis Africana Oil on The Chicken Breast Flesh

Fatty Acids	T1	T2	T3	T4	T5	T6	SEM	P-value
C14:0	3.10 ^a	2.91 ^b	2.41 ^b	2.21 ^b	2.19 ^c	2.02 ^c	0.63	0.08
C16:0	22.1 ^a	20.0 ^b	14.2 ^b	14.1 ^c	13.7 ^c	12.7 ^c	1.30	0.02
C18:0	11.1 ^a	10.3 ^b	6.72 ^b	6.01 ^c	5.64 ^c	5.02 ^c	0.72	0.80
C20:0	4.10 ^a	3.11 ^a	2.50 ^b	2.26 ^b	1.81 ^c	1.73 ^c	0.23	0.08
C22:0	0.30 ^a	0.27 ^b	0.20 ^b	0.15 ^b	0.14 ^b	0.10 ^b	0.04	0.29
C14:1c	1.25 ^c	1.14 ^a	2.91 ^a	3.11 ^a	3.90 ^a	3.94 ^a	0.08	0.46
C16:1c	2.01 ^b	2.71 ^b	3.41 ^b	3.48 ^a	3.56 ^a	4.03 ^a	0.06	0.50
C18:1c	10.9 ^c	14.2 ^b	20.1 ^b	21.8 ^a	23.1 ^a	23.71 ^a	1.05	0.40
C18:1n9t	1.00 ^c	1.13 ^b	1.51 ^b	1.54 ^b	1.80 ^a	1.93 ^a	0.08	0.27
C18:1n9c	0.61	0.80	1.00	1.05	1.40	1.44 ^a	0.03	0.21
C:22:1	0.10 ^b	0.17 ^a	0.84 ^a	0.92 ^a	1.00 ^a	1.02 ^a	0.06	0.61
C18:2n6	11.7 ^c	14.0 ^b	21.5 ^a	23.1 ^a	24.6 ^a	25.1 ^a	0.54	0.37
C18:5n3	0.61 ^c	0.65 ^b	0.93 ^b	1.10 ^b	1.28 ^a	1.34 ^a	0.07	0.75
C18:3n3	2.80 ^c	3.09 ^b	7.36 ^b	9.19 ^a	11.3 ^a	13.9 ^a	0.40	0.31
C20:4n6	1.17 ^b	1.85 ^b	2.50 ^b	2.67 ^b	3.06 ^a	3.21 ^a	0.18	0.20
C20:3n6	0.42 ^b	0.78 ^a	1.10 ^a	1.35 ^a	1.40 ^a	1.64 ^a	0.12	0.004
C22:6n3	0.10 ^c	0.25 ^b	0.84 ^b	1.02 ^b	1.63 ^a	1.72 ^a	0.10	0.02
TSFA ¹	53.80 ^a	50.70 ^a	39.21 ^b	18.67 ^c	17.80 ^c	16.94 ^c	2.71	0.21
TUFA ²	36.20 ^c	49.30 ^b	79.45 ^a	81.33 ^a	82.20 ^a	83.06 ^a	4.47	0.05

MUFA ³	20.08 ^c	27.90 ^b	34.10 ^a	35.22 ^a	35.60 ^a	36.06 ^a	2.32	0.01
PUFA ⁴	26.12 ^c	31.40 ^b	45.35 ^a	46.11 ^a	46.60 ^a	47.00 ^a	2.60	0.02
n-6:n-3 ⁵	3.58 ^a	3.11 ^a	3.00 ^a	2.92 ^b	2.77 ^b	2.60 ^b	0.03	0.01
Ant. Index ⁶	0.92 ^a	0.97 ^a	0.62 ^b	0.50 ^b	0.36 ^c	0.25 ^c	0.01	0.001

¹Total saturated fatty acid= C14:0 + C16:0 + C18:0 + C20:0 +C22:0

²Unsaturated fatty acid [TSFA] = (3 + 4)

³Mono unsaturated fatty acid [TMUFA] = C14:1C + C16:1C + C18:1C + C18:1n9t + C18:1n9c + C22:1

⁴Polyunsaturated fatty acid [TPUFA] = C18:2 n6 + C20:5 n3 + C18:3n3 + C20:4n6 + C20:3n6 + C: 22:6n3

⁵n-6: n-3 = (C18:2 n6 + C20:4n 6 + C20:3n 6 / (C20:5n 3 + C18:3n 3 + C:22 6n 3), ⁶Antherogenic index = (C12:0+ 4×C14:0+ C16)/Σ of UFA

Potential Impact of Prosopis Africana Oil on The Chicken Broiler's Thigh Muscle

The impact of Prosopis africana oil here on thigh muscle of broiler chickens is shown in Table 4. Myristic acid [C14:0; 2.18 – 3.18 percent], palmitic acid [C16:0; 14.3 – 22.9 percent], stearic acid [C18:0; 4.18 – 11.20 percent], arachidic acid [C:20; 1.45 – 4.21 percent], behenic acid [C:22; 0.20 – 0.31 %], myristoleic acid [C14:1c; 1.81 – 2.66 %], palmitoleic acid [C16:1c; 2.01 – 3.40 percent], oleic acid [C18:1n 9t; 1.20 – 2.00 percent], elaidic acid [C18:1n 9c; 0.51 – 1.00 %], erucic acid [C22:1; 0.13 – 0.87 percent], vaccenic acid [C18:2n6; 15.4 – 24.0 percent], linolenic acid [C18:5n3; 0.88 – 1.89 percent], α-linolenic acid [C18:3n3; 3.04 – 15.6 percent], eicosapentaenoic acid [C20:4n6; 2.08 – 4.00 percent], docosahexaenoic acid [C20:3n6; 0.92 – 1.88 percent] and linoeladic acid [C22:6n3; 0.05 – 2.31 percent], total saturated fatty acid [TSFA; 20.18 – 49.80 percent], monosaturated fatty acid [TMUFA; 23.12 – 31.22 percent] and polyunsaturated fatty acid [TPUFA; 27.08 – 47.60 percent]. The highest levels of PUFA and MUFA were found in T3-T6, whereas the lowest levels were found in T1 (P<0.05). The therapies had a substantial (P<0.05) impact on TSFA.

Table 4: Potential Impact of Prosopis Africana Oil on The Chicken Broiler's Thigh Muscle

F. Acids	T1	T2	T3	T4	T5	T6	SEM	P-value
C14:0	3.18 ^a	2.48 ^b	2.42 ^b	2.37 ^b	2.21 ^c	2.18 ^c	0.08	0.32
C16:0	22.9 ^a	18.0 ^b	17.2 ^b	16.3 ^c	15.0 ^c	14.3 ^c	0.51	0.61
C18:0	11.2 ^a	8.71 ^b	7.62 ^b	6.00 ^c	5.73 ^c	4.18 ^c	0.30	0.31
C20:0	4.21 ^a	3.93 ^a	2.81 ^b	2.40 ^b	1.88 ^c	1.45 ^c	0.73	0.74
C22:0	0.31 ^a	0.21 ^b	0.27 ^b	0.20 ^b	0.24 ^b	0.20 ^b	0.07	0.005
C14:1c	1.81 ^c	2.88 ^a	2.73 ^a	2.91 ^a	2.97 ^a	2.66 ^a	0.08	0.007
C16:1c	2.01 ^b	2.21 ^b	2.93 ^b	3.18 ^a	3.51 ^a	3.40 ^a	0.04	0.28
C18:1c	13.4 ^c	18.7 ^b	19.5 ^b	21.0 ^a	21.7 ^a	22.8 ^a	0.26	0.02
C18:1n9t	1.20 ^c	1.51 ^b	1.40 ^b	1.44 ^b	1.86 ^a	2.00 ^a	0.06	0.02
C18:1n9c	0.51 ^c	0.56 ^c	0.80 ^b	0.83 ^b	0.87 ^b	1.00 ^a	0.02	0.11
C:22:1	0.13 ^b	0.47 ^a	0.51 ^a	0.53 ^a	0.66 ^a	0.87 ^a	0.04	0.64
C18:2n6	15.4 ^c	19.8 ^b	22.0 ^a	22.8 ^a	23.4 ^a	24.0 ^a	0.23	0.16
C20:5n3	0.88 ^c	1.05 ^b	1.10 ^b	1.14 ^b	1.51 ^a	1.89 ^a	0.18	0.0001
C18:3n3	3.04 ^c	8.08 ^b	10.3 ^b	13.1 ^a	14.3 ^a	15.6 ^a	0.22	0.06
C20:4n6	2.08 ^b	2.21 ^b	2.59 ^b	2.87 ^b	3.81 ^a	4.00 ^a	0.05	0.02
C20:3n6	0.92 ^b	1.08 ^a	1.17 ^a	1.21 ^a	1.28 ^a	1.88 ^a	0.01	0.001
C22:6n3	0.05 ^c	1.35 ^b	1.40 ^b	1.44 ^b	2.00 ^a	2.31 ^a	0.61	0.03
TSFA1	49.80 ^a	40.56 ^a	30.40 ^b	25.80 ^c	21.56 ^c	20.18 ^c	0.06	0.05
TUFA2	50.20 ^c	56.14 ^b	70.60 ^a	75.20 ^a	79.44 ^a	78.82 ^a	0.02	0.001
MUFA3	23.12 ^c	25.70 ^b	29.85 ^b	31.41 ^a	32.38 ^a	31.22 ^a	0.14	0.26
PUFA4	27.08 ^c	33.74 ^b	40.75 ^a	43.79 ^a	47.06 ^a	47.60 ^a	0.47	0.02
n-6:n-35	3.91 ^a	3.00 ^a	2.97 ^b	2.80 ^b	2.77 ^b	2.59 ^b	0.03	0.001
Ant. Ind6	0.82 ^a	0.72 ^a	0.54 ^b	0.50 ^b	0.42 ^c	0.38 ^c	0.05	0.012

¹Total saturated fatty acid [TSFA] = C12:0 + C14:0 + C16:0 + C18:0 + C20:0 +C22:0

²Unsaturated fatty acid = (3 + 4)

³Mono unsaturated fatty acid [TMUFA] = C14:1C + C16:1C + C18:1C + C18:1n9t + C18:1n9c + C22:1

⁴Polyunsaturated fatty acid [TPUFA] = C18:2 n6 + C20:5 n3 + C18:3n3 + C20:4n6 + C20:3n6 + C: 22:6n3

⁵n-6: n-3 = (C18:2 n6 + C20:4n 6 + C20:3n 6 / (C20:5n 3 + C18:3n 3 + C:22 6n 3),

⁶Antherogenic index = (C12:0+ 4×C14:0+ C16)/Σ of UFA

Sensory Assessment of Fed Broiler Chicken Fed Prosopis Africana Oil

Table 5 presents the sensory evaluation of broiler chicken given various amounts of Prosopis africana oil. The color, flavor, texture, juiciness, tenderness, and general acceptability of the breast and thigh meat were all significantly ($P<0.05$) influenced by the treatments.

Table 5: Sensory Assessment of Broiler Chicken Fed Prosopis Africana

Parameters	T1	T2	T3	T4	T5	T6	SEM	P-value
Breast meat								
Colour	4.62 ^b	4.90 ^b	6.01 ^a	6.50 ^a	6.59 ^a	6.73 ^a	0.10	0.96
Flavour	5.05 ^b	5.20 ^b	6.18 ^a	6.22 ^a	6.27 ^a	6.35 ^a	0.09	0.60
Texture	4.73 ^b	4.86 ^b	5.40 ^a	5.51 ^a	5.60 ^a	5.62 ^a	0.08	0.58
Juiciness	5.06 ^c	5.11 ^c	6.86 ^b	6.92 ^b	7.06 ^a	7.11 ^a	0.11	0.72
Tenderness	4.10 ^c	4.42 ^c	5.92 ^b	6.06 ^a	6.12 ^a	6.19 ^a	0.10	0.69
Acceptability	4.96 ^d	5.00 ^c	6.67 ^b	6.91 ^b	6.96 ^b	7.60 ^a	0.07	0.50
Thigh meat								
Colour	4.09 ^c	4.62 ^c	6.61 ^b	6.68 ^b	7.03 ^a	7.05 ^a	0.24	0.07
Flavour	3.81 ^b	3.90 ^b	6.00 ^a	6.04 ^a	6.81 ^a	6.88 ^a	0.06	0.17
Texture	5.06 ^b	5.11 ^b	7.03 ^a	7.30 ^a	7.50 ^a	7.55 ^a	0.30	0.57
Juiciness	3.60 ^c	4.57 ^b	6.29 ^b	7.00 ^a	7.08 ^a	7.15 ^a	0.04	0.72
Tenderness	4.71 ^d	5.33 ^c	6.92 ^b	7.00 ^a	7.08 ^a	7.21 ^a	0.94	0.59
Acceptability	4.29 ^d	5.86 ^c	6.22 ^b	6.83 ^b	6.90 ^b	7.10 ^a	0.05	0.87

Standard error of the mean (SEM) T1: control diet with no Prosopis africana oil; T2: control diet plus 2.5g/kg oxytetracycline; T3: baseline diet plus 200 mg; T4: baseline diet plus 400 mg; T5: control diet plus 600 mg; T6: basal diet plus 800 mg Prosopis africana oil

Discussion

Poultry meat has a low lipid content and a disproportionately high quantity of polyunsaturated fatty acids, according to [29, 30,39]. The amount and make-up of meat fat may shift in response to alterations in the lipid profile of an animal's diet [30-34]. The results of the present investigation showed that, in comparison to the other treatments, dietary supplementation of Prosopis africana oil enhanced the contents of polyunsaturated fatty acids (TPUFA) in the breast and thigh meat ($P<0.05$). Saturated fatty acid (TSFA) on the other hand, declined as Prosopis africana oil levels increased in all regimens. The current finding is consistent with those of [35-37]. who found that broiler chicken meat given two herbal extract and virginiamycin had increased TPUFA and reduced TSFA contents. Similar findings were made by, who found that broiler chicken thigh meat had larger TPUFA concentrations and a better n-6: n-3 fatty acid ratio when oregano essential oil was added to the diet at a rate of 400 mg/kg [38,40]. These results provide strong evidence that Prosopis africana oil can alter the nutritional value of broiler meat because phytonutrients were present in the sample. Assert that appropriate antibody synthesis in animals depends on the dietary n-6: n-3 ratio [41-43]. To prevent inflammatory reactions and other associated health risks, the ratio of omega-3 to omega-6 fatty acids ought to be within 1 and 4 [44, 46]. In this investigation, broiler chicks given Prosopis africana oil showed a more favorable ratio of omega-6 to omega-3 fatty acids. Dietary Prosopis africana oil was found to alter monosaturated fatty acids (TMUFA) in the flesh, it increases more noticeably in the breast than the thigh, which results in a larger overall concentration of TMUFA. Noted an increase in TMUFA in the breast meat of broilers given green tea extract, which is consistent with the current findings [45].

Additionally, broiler chickens given various combinations of essential oils showed a higher concentration of MUFA in the flesh of their breast and thighs. According to [46]. The presence of phytonutrients in Prosopis africana oil are helpful in lowering

the surplus synthesizing TSFA, which inhibits the biosynthesis of TMUFA, may be responsible for the rise in TMUFA contents in the breast and thigh meat [45]. A rise in high density lipoprotein (HDL) will lead to arise in the ratio of HDL cholesterol to overall cholesterol as well as the Anthrogenicity index. Reported that antherogenicity index (ANI) is used to gauge the safety of meat; the lower the ANI, the more secure the meat is, and vice versa. This suggests that essential oils have no adverse effects on the lipid quality [28,30].

Evaluation of the Sensory Assessment in Broiler Chicken Fed Prosopis Africana Oil

The present study's findings revealed that the cooked breast and thigh meat had considerable ($P<0.05$) differences in color, flavor, tenderness, juiciness, texture, and general acceptability. The current observations are consistent with those of [47] who discovered that feeding broiler chickens 200 mg/kg of clove and lemongrass oil considerably ($P<0.05$) changed how the meat's flavor was perceived. [48] found comparable outcomes, noting that the addition of thyme oil can significantly alter the taste, softness, color, and general acceptance of the meat between groups. It was discovered that the chicken flesh samples given 300 mg/kg thyme oil were tastier, softer, and had a more vibrant color. Because the panelists judged that the thigh and breast meat from broilers administered Prosopis africana oil had better texture, flavor, tenderness, juiciness, and general acceptance relative to other treatments, the results of this trial imply that the addition of Prosopis africana oil could improve taste of broilers meat.

Conversely, reported that dietary supplementation of garlic oil at 50 mg/kg did not significantly ($P>0.05$) alter the color, softness, flavor, or general acceptance of breast meat [49]. The disparities in these results could be attributed to the differences in the composition of bioactive compounds in essential oils. The occurrence of carotene in Prosopis africana oil may have impacted the substantial variations in color found ($P<0.05$) in the breast and

thigh meat [28]. The current condition of the meat colors, pre-slaughter variables (genetics, feed, managing, stress, temperatures and cold stress, gaseous surroundings), slaughter, chilling, and subsequent processing situations (stunning methods, inclusion of nitrates, components, pH, last cooking temperatures, reducing circumstances, and irradiation) are the main factors affecting meat color [50-53]. The maturation of the connective tissues, the contractile condition of the myofibrillar proteins, environmental stress, scalding temperature, the maturity level of the birds, the rate of rigor improvement, the rate of cooling, and the filleting period are the primary variables impacting meat tenderness [51,54].

Conclusion

It was determined that Prosopis africana oil has a number of phyto-constituents with the potential to alter the fatty acid composition of meat. Its addition to the diets of birds is not linked to any adverse consequences, such as the deposit of hazardous carcass residue that are related to with the use of antibiotic growth promoters. Broilers' diet can be supplemented with Prosopis essential oil (800 mg/kg) to enhance sensory attributes.

References

1. Caroline CV (2019) Improving feed efficiency and gut integrity in Escherichia coli challenged weaned pigs. International Pig Magazine 1: 8-10.
2. Sandra C (2020) Phyto-genics: how to improve pig production efficiency with plants. International Pig Magazine 8: 1-3.
3. Chris, C (2010) Gut flora and immune modulation by plant extracts. International Poultry Magazine 7: 1-2.
4. Man DL (2006) Support gut health with a unique combination of fatty acid and phyto-genics. International Poultry Magazine 7: 6-8.
5. Jan D, Van K, and Ester, V (2016) Phyto-genics: be one step ahead with plant derived feed additives. International Poultry Magazine 5: 3-5.
6. Alagbe, Olujimi John, Oluchi CP Agubosi and Rufus et al. (2023) Histopathology of broiler chickens fed diet supplemented with Prosopis africana (African mesquite) essential oil. Brazilian Journal of Science 2: 49-59.
7. Alagbe, Olujimi John, Oluchi C Precious Agubosi, Rufus et al. (2023) Haemato-biochemical indices and intestinal microbial population of broiler chickens fed diet supplemented with Prosopis africana (African mesquite) essential oil. Brazilian Journal of Science 2: 98-110.
8. Alagbe JO, Bamigboye, S, Nwosu GC, Agbonika DA and Kadiri Mercy Cincinsoko Et al. (2023) Characterization of bioactive compounds in Luffa aegyptiaca leaf ethanolic extracts using gas chromatography and mass spectrometry (GC-MS). Drug Discovery 17: e10dd1011
9. Alagbe JO, Kadiri MC, Oluwafemi RA, Agubosi OCP and Anorue DN et al. (2023) Analysis of bioactive compounds in ethanolic extracts of Xylopia aethiopica leaves using gas chromatography and mass spectrometry technique. American Journal of Science on Integration and Human Development 1: 1-10.
10. Alagbe JO (2023) Bioactive compounds in ethanolic extract of Strychnos innocua root using gas chromatography and mass spectrometry (GC-MS). Drug Discovery 17:e4dd1005.
11. Alagbe John Olujimi, Ramalan Sadiq Muhammad., Shittu Muritala Daniel and Olagoke Olayemi Christiana (2022) Effect of Trichilia monadelpha stem bark extract on the fatty acid composition of rabbit's thigh meat. Journal of Environmental Issues and Climate Change 1: 63-71
12. Alagbe JO, Shittu MD and Tanimomo, Babatunde K (2022) Influence of Anogeissus leio carpus stem bark on the fatty acid composition in meat of broiler chickens. European Journal of Life Safety and Stability 14: 13-22.
13. Alagbe JO (2022) Use of medicinal plants as a panacea to poultry production and food security: A review. Gospodarka I Innowacje 22: 1-12.
14. Singh Sharma, Alagbe Olujimi John, Liu Xing, Sharma Ram and Kumar Amita (2022) Comparative analysis of ethanolic Juniperus thurifera leaf, stem bark and root extract using gas chromatography and mass spectroemetry. International Journal of Agriculture and Animal Production, 2: 18-27.
15. Agubosi OCP, Alexander, James and Alagbe JO et al. (2022) Influence of dietary inclusion of Sunflower (Helianthus annus) oil on growth performance and oxidative status of broiler chicks. Central Asian Journal of Medical and Natural Sciences 2: 187-195.
16. Agubosi OCP, Soliu MB and Alagbe JO (2022) Effect of dietary inclusion levels of Moringa oleifera oil on the growth performance and nutrient retention of broiler starter chicks. Central Asian Journal of Theoretical and Applied Sciences 3: 30-39.
17. Agubosi OCP, Imudia, Favour Dumkenechukwu and Alagbe JO et al. (2022) Evaluation of the nutritional value of air dried and sun-dried sweet potato (Ipomoea batatas) peels. European Journal of Life Safety and Stability 14: 43-51.
18. Alagbe JO (2022) Gas chromatography and mass spectroscopy of Juniperus phoenice stem bark extract and its influence on the haemato-biochemical values of growing rabbits. British Scientific Periodical 1: 18-33.
19. Alagbe JO, Zubairu Habiba, Adedeji OM, Bamigboye S and Dora Agbonika et al. (2022) Influence of Juniperus thurifera root extract on the nutrient digestibility and caecal microbial count of growing rabbits. Web of Synergy: International Interdisciplinary Research Journal 1: 5-17.
20. Alagbe JO, Shittu MD, Ramalan SN, Tanimomo KB and Adekunle D et al. (2022) Growth performance, semen quality characteristics and hormonal profile of male rabbit bucks fed Rubia cordifolia root extracts. International Journal of Biological Engineering and Agriculture 1: 1-13.
21. Alagbe JO, Agubosi OCP, Oluwafemi RA and Gabriel Zakara et al. (2022) Efficacy of Trichilia monadelpha stem bark on the growth performance of growing rabbits. Best Journal of Innovation in Science, Research and Development 1: 10-19.
22. Shittu MD, Alagbe JO, Rafiu TA, Oye KE, Oladokun PO and Agbaje IA et al. (2020) Physiological response of mixed breed rabbits fed varied replacement levels of maize with cassava root meal. European Journal of Biotechnology and Bioscience, 8: 16-21.
23. Alagbe JO and Akintayo-Balogun OM (2020) Effects of dietary supplementation of Albizia lebbek seed oil (ALO) on the fatty acid composition of weaner rabbits. Biochemistry and Biotechnology Research 8: 29-33.
24. Shittu, M.D and Alagbe, J.O (2020) Phyto-nutritional profiles of broom weed (Sida acuta) leaf extract. International Journal of Integrated Education. 3: 119-124
25. Akintayo Balogun Omolere M and Alagbe JO (2020) Probiotics and medicinal plants in poultry nutrition: A review. United International Journal for Research and Technology, 2: 7-13.
26. Alagbe JO (2020) Caecal Microbial Population of Growing Grass Cutters (Thyronoymys Swinderianus) Fed Phyllanthus Amarus and Pillogstigma Thonnigii Leaf Meal Mixture as Partial Replacement for Soya Bean Meal. Concept of Dairy and Veterinary Sciences 3: 350-355.

27. Oluwafemi RA, Oluwayinka EO and Alagbe JO (2020) Effect of dietary supplementation of neem oil (*Azadirachta indica*) on the growth performance and nutrient digestibility of weaned rabbits. *European Journal of Biotechnology and Bioscience*. 8: 6-10.
28. Alagbe JO, Adeoye, Adekemi and Oluwatobi OA (2020) Proximate and mineral analysis of *Delonix regia* leaves and roots. *International Journal on Integrated Education*. 3: 144-149.
29. Alagbe JO, Shittu MD and Tanimomo, Babatunde K (2022) Influence of *Anogeissus* stem bark on the fatty acid composition in meat of broiler chickens. *European Journal of Life Safety and Stability* 14: 13-22.
30. Adewale AO, Alagbe JO, Adeoye, Adekemi O (2021) Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phyto-genic Feed Additive in Growing Rabbit Diets: Haematology and serum biochemical indices. *International Journal of Orange Technologies* 3: 1-12
31. Al-Kassie GAM (2009) Influence of two plant extracts derived from thyme and cinnamon on broiler performance. *Pakistan Veterinary Journal* 29:169-173.
32. Alagawany M, Ashour EA and Feda FM (2015) Effect of dietary supplementation of garlic (*Allium sativum*) and turmeric (*Curcuma longa*) on the performance, carcass traits, blood profile and oxidative status in growing rabbits. *Annals of Animal Science*, 16: 489-505.
33. Erener GN, Ocak A, Altop S, Cankaya H, Murat A and Ozturk E et al. (2011) Growth performance, meat quality and caecal microbial count of broilers fed green tea extract. *Asian-Australian Journal of Animal Science* 24: 1128-1135.
34. Hashemipour H, Kermanshahi H, Golian A and Veldkamp T (2013) Effect of thyme feed supplementation on the performance, antioxidant enzyme activities and fatty acid composition in broiler chickens. *Poultry Science* 2: 2059-2069.
35. Camy MLY, Redoy MRA, Shuvo AAS, Rahman MA and Al-Mamum M (2019) Effect of aqueous herbal extract on growth, plasma metabolites and meat characteristics of broilers. *Bangladesh Journal of Animal Science* 48; 108-115.
36. Yang JE, Ye Seul Seo, Muhammad AD, Hong SM, Hyeoung SP and Chul JY et al. (2020) Physico-chemical attributes sensory evaluation and oxidative stability of leg meat from broilers supplemented with plant extracts. *Journal of Animal Science and Technology* 62: 730-740.
37. Mohammed HN (2013) Study on some chemical, physical, sensory and bacteriology characteristics of canned chicken meat imported in Sulaymaniyah markets. *Iraq International Nutrition* 5: 128-133.
38. Puvaca N, Kostadinovic L, Popovic S, Ljubojevic D and Tufarelli V et al. (2015) Proximate composition, cholesterol concentration and lipid oxidation of meat from chickens fed dietary spice addition (*Allium sativum*, *Piper nigrum*, *Capsicum annum*). *Animal Production Science* 56: 1920-1927.
39. Jang A, Liu X, Shin M, Lee B, Lee S and Lee J et al. (2008) Antioxidative potential of raw breast meat from broiler chicks fed a dietary medicinal herb extract mix. *Poultry Science* 13: 124-136.
40. Florou PP, Palatos G, Govaris D, Botsoglou I, Giannenas I and Ambrosiadis A et al. (2005) Oregano herb versus oregano essential oil as feed supplementation to increase the oxidative stability in poultry meat. *International Journal of Poultry Science* 4: 866-871.
41. Purohit A and Vyas KB (2006) Antiatherosclerotic effect of *Capparis decidua* fruit extract in cholesterol-fed rabbits. *Pharmacology* 44: 172-177.
42. Mohammed AT, Serkan Y and Mehmet AY (2019) Effects of tarragon (*Artemisia dracunculus*) powder on broiler performance parameters and histopathology of internal organs. *Australian Journal of Veterinary Science* 51: 113-118.
43. Malahubban M and Aziz B (2016) Serum biochemical properties and liver morphology of broiler chicken affected by feeding Masai kucing as a supplementary diet. *Journal of Research and Biotechnology* 8: 123-131.
44. Sameh A, Mahmoud A, Mohammed EA, Asmaa MS, Islam MS et al. (2018) Growth, carcass traits, blood haematology, serum metabolites, immunity and oxidative indices of growing rabbits fed diets supplemented with red or black pepper oils. *Animals*, 8: 168-173.
45. Yesilbag D, Eren M, Agel H, Kovanlikaya A (2011) Effect of dietary rosemary oil and vitamin E on performance, meat quality and serum SOD activity. *British Journal of Poultry Science* 52: 4722-4826.
46. Emami NK, Samie A, Rahmani HR and Ruiz-Feria CA (2012) The effect of peppermint essential oil and fructooligosaccharides as alternatives to virginiamycin on growth performance, digestibility, gut morphology and immune response of male broilers. *Animal Feed Science and Technology* 175: 57-64.
47. Ilias G, Eleftherios B, Efterpi C and Panagiota FP (2013) Essential oils and their applications in animal nutrition. *Medicinal and Aromatic Plants* 2: 2-12.
48. Krishan G and Narang A (2014) Use of essential oil in poultry nutrition: A new approach. *Journal of Veterinary and Animal Research* 1: 156-162.
49. Reda FM, Alagawany M, Mahmoud HK, Mahgoub SA (2020) Use of red pepper oil in quail diets and its effect on performance, carcass measurements, intestinal microbiota, antioxidant indices, immunity and blood constituents. *Animal* 14: 1025-1033.
50. Sarica S, Corduk M, Yarim GF, Yenisehirli G (2009) Effects of novel feed additives in wheat-based diets on performance carcass and intestinal tract characteristics of quail. *South African Journal of Animal Science* 39: 144-157.
51. Torki M, Sedgh-Gooya S and Mohammadi H (2018) Effects of adding essential oil of rosemary and chicory extract to diets on performance, egg quality and some blood parameters of laying hens subjected to heat stress. *Journal of Applied Animal Research* 46: 1118-1126.
52. Berri C, Wacrenier N and Millet N (2001) Effect of selection for improved body weight and meat characteristics of broilers from experimental and commercial lines. *Poultry Science* 80 : 833-838.
53. Broncano JM, Petron MJ, Parr V and Timon ML (2009) Effect of different cooking methods on lipid oxidation and formation of free cholesterol oxidation products in *Latissimus dorsi* muscle of Iberian pigs. *Meat Science* 83: 431-437.

Copyright: ©2023 Alagbe. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.