

Case Report

Open Access

Metastatic Urothelial Carcinoma of Occult Primary Origin Revealed by a Necrotic Inguinal Lymph Node: A Rare Case Report

Bjane O*, Taïbou T, Tmiri A, El Badr M, Deghdagh Y, Kbiro A, Moataz A, Dakir M, Debbagh A and Aboutaieb R

Urology Department, Ibn Rochd Hospital, Faculty of Medicine and Pharmacy, Hassan II University, Casablanca, Morocco

ABSTRACT

Introduction: Occult primary urothelial carcinoma is an extremely rare entity that may present as isolated lymph node involvement without visible bladder or ureteral lesions, making diagnosis particularly challenging.

Case Presentation: We report the case of a 91-year-old man presenting with a left inguinal swelling evolving over four years. Examination revealed a necrotic, fistulized mass with no urothelial lesions on cystoscopy. Biopsy showed a poorly differentiated carcinoma (CK7+/CK20+/p63+/p40+/GATA3-), initially suggesting squamous differentiation, but the lymphatic distribution and suspicious ureteral thickening supported an occult urothelial origin. A multidisciplinary management approach was undertaken. Systemic chemotherapy was not initiated due to age and comorbidities. Local wound care and antibiotic therapy led to partial improvement and clinical stabilization without visceral progression.

Discussion: This case illustrates the diagnostic complexity of dedifferentiated urothelial carcinomas with atypical immunohistochemical profiles. Loss of GATA3 expression, though uncommon, does not exclude urothelial origin, especially in high-grade tumors. Presentation as isolated inguinal lymphadenopathy is exceedingly rare.

Conclusion: Diagnosis of occult urothelial carcinoma requires multidisciplinary collaboration and careful integration of clinical, radiological, and pathological data. In elderly or frail patients, a conservative, symptom-oriented approach is often the most appropriate. Reporting such rare cases is essential to improve understanding of the atypical presentations and dissemination pathways of urothelial carcinoma.

*Corresponding author

Bjane O, Urology Department, Ibn Rochd Hospital, Faculty of Medicine and Pharmacy, Hassan II University, Casablanca, Morocco.

Received: October 24, 2025; **Accepted:** October 28, 2025; **Published:** November 07, 2025

Keywords: Urothelial Carcinoma, Occult Primary, Inguinal Lymphadenopathy, GATA3, Immunohistochemistry, Case Report

Introduction

In cancer diagnosis and management, accurately identifying the primary origin of tumor cells is crucial, as it directly guides therapeutic decisions. Occult cancer refers to a rare condition in which the primary lesion remains clinically undetectable, making diagnosis challenging. In such cases, the metastatic lesion is typically identified first through clinical evaluation and histopathological examination. Occasionally, the microscopic primary tumor is only discovered post-mortem during autopsy examinations [1].

Most reported cases of occult malignancy involve breast cancer, with fewer instances observed in thyroid and gynecologic origins [1-2]. Rarely, certain occult somatic tumors may initially present as acute cerebral infarction [3,4]. The incidence of occult cancer remains very low, representing only 0.3% to 1.0% of newly diagnosed breast cancers [5,6].

Urothelial carcinoma (transitional cell carcinoma) arises from the transitional epithelium of the urinary tract. Although it typically

develops within the urinary bladder, ureter, or renal pelvis, distant metastases are uncommon and usually appear only in advanced disease stages following diagnosis. To date, presentation of urothelial carcinoma with metastasis as the initial clinical manifestation has been exceedingly rare.

Case Presentation

A 91-year-old male with no significant surgical history had a past medical history notable for tuberculous pericarditis successfully treated five years earlier.

The patient's history dated back approximately four years, marked by the gradual onset of a painless left inguinal swelling, associated with edema of the left lower limb. The course was slowly progressive, with a gradual increase in tumor size over the past two years, without fever or pain, evolving in a context of apyrexia and relatively preserved general condition.

On physical examination, the patient was alert, in good general condition, with a Performance Status (PS) of 1. The conjunctivae were normally colored, and edema of the left lower limb was noted. Urological examination showed a preserved diuresis, without

lumbar tenderness or suprapubic pain. Locally, there was a left inguinal swelling fistulized to the skin, firm and painless, without acute inflammatory signs.

The external genitalia appeared normal. Digital rectal examination revealed a prostate estimated at 50 grams, firm and regular, with a supple bladder base.

Biological investigations showed hemoglobin at 12.7 g/dL, C-Reactive Protein (CRP) at 184 mg/L, and leukocytosis at 14,310/ μ L, indicating a biological inflammatory syndrome.

Renal function was preserved (creatinine 12.7 mg/L, urea 0.35 g/L) and serum potassium was normal (4.58 mEq/L). Urine culture was sterile, and total PSA was 3.14 ng/mL, within normal limits for the patient's age.

Inguinal ultrasound performed in October 2021 revealed bilateral hypogastric and left inguinal lymphadenopathy, some with suspicious features.

Thoraco-abdominopelvic CT (TAP) demonstrated a dilated and tortuous lumbar ureter, upstream of a parietal thickening at the L5–S1 level, measuring approximately 36 mm and enhancing after contrast injection—suggestive of a possible ureteral infiltrating lesion.

The urinary bladder contained a right lateral wall diverticulum without suspicious thickening or mucosal irregularity, while the prostate was enlarged (66.8 mL).

At the left inguinal region, there was a heterogeneous, ulcerative-necrotic mass, poorly demarcated, with heterogeneous post-contrast enhancement and central necrotic areas. The mass encased the superficial femoral artery and vein (which remained patent), infiltrated the sartorius muscle laterally and the adductor longus medially, and was associated with diffuse infiltration of adjacent subcutaneous fat.

Bilateral iliac lymphadenopathy was also noted, more pronounced on the left side (largest node measuring 24.9×20.5 mm).



Figure 1: Left Inguinal Mass Fistulized to the Skin, Showing a Necrotic Ulceration with Seropurulent Exudate



Figure 2: Evolution After Three Months: Drier Ulcerated Necrotic Wound with Reduced Exudation, Indicating Partial Local Improvement

At the thoracic level, there was circumferential pleural thickening of the right hemithorax and three hepatic arterial phase enhancements.

Cystoscopy showed a normal urethra, two visible ureteral orifices, and a bladder of good capacity, with a diverticulum of the right lateral wall and bladder base, but no mucosal lesions or suspicious thickening.

Biopsy of the left inguinal mass revealed a secondary subcutaneous localization of a poorly differentiated carcinoma.

Immunohistochemical (IHC) staining demonstrated a CK7+/CK20+, p63+, p40+, and GATA3– profile, initially suggestive of a squamous cell carcinoma.

However, several morphological and clinical findings—notably the enhancing ureteral thickening, the presence of a thick-walled bladder diverticulum, and the pelvic and inguinal nodal distribution—strongly supported a possible occult urothelial origin.

Although GATA3 negativity is more typical of squamous differentiation, it does not exclude urothelial carcinoma, particularly in high-grade dedifferentiated tumors, where GATA3 loss has been reported in up to 10–20% of cases.

The absence of any identifiable cutaneous, rectal, prostatic, or digestive primary lesion further strengthened the hypothesis of a high-grade metastatic urothelial carcinoma of unknown primary origin.

PET-CT revealed hypermetabolic superficial and femoral left inguinal lymph nodes, consistent with secondary involvement, and a right bladder diverticulum with a thickened wall requiring further evaluation.

It also showed diffuse interstitial pneumonia with mediastinal hypermetabolic lymph nodes, likely inflammatory or infectious, and no other suspicious foci of hypermetabolism, particularly in the soft tissues of the lower limbs.

Follow-up CT TAP confirmed the persistence of necrotic left inguinal lymph nodes, with increase in size and persistent cutaneous fistulization.

The posterolateral bladder diverticulum remained unchanged and non-suspicious, and hepatic cystic lesions corresponded to simple biliary cysts. Finally, colonoscopy demonstrated a normal rectocolonic mucosa with no evidence of malignancy.

Taken together, the morphological, radiological, and immunohistochemical findings a CK7+/CK20+, p63/p40+ carcinoma, associated with pelvic and inguinal lymph node metastases, suspicious ureteral thickening, and a bladder diverticulum with wall irregularity favored the diagnosis of a metastatic urothelial carcinoma of occult primary origin.

This case illustrates the rarity and diagnostic complexity of dedifferentiated urothelial carcinomas, which may mimic squamous cell carcinoma and initially present with isolated inguinal lymphadenopathy.

A multidisciplinary management approach was undertaken, involving the urology, medical oncology, and pathology teams.

Locally, control of the inguinal mass initially relied on daily local wound care, including cleansing of the cutaneous fistula, application of absorbent and antiseptic dressings, and infection prevention.

A probabilistic antibiotic therapy was initiated because of the infectious risk related to tumor necrosis and cutaneous fistulization, resulting in favorable clinical evolution and progressive drying of the wound.

On a systemic level, after multidisciplinary discussion, systemic chemotherapy was not initiated due to the expected tolerance profile in a very elderly patient and the high risk of hematologic and renal toxicity.

An oral chemotherapy regimen or carboplatin-based treatment was considered but not implemented, given the frailty of the patient and borderline renal function.

Immunotherapy, which is recommended in locally advanced or metastatic urothelial carcinomas not eligible for platinum-based chemotherapy, was also discussed.

However, due to the absence of a histologically confirmed primary urothelial lesion in the bladder or ureter, and uncertainty regarding the exact tumor origin, this option was not pursued.

Radiological follow-up showed persistent necrotic left inguinal lymphadenopathies, with an increase in size and persistent cutaneous fistulization, but no evidence of new distant metastases.

Clinically, the patient remained stable over several months, with preserved diuresis, absence of pelvic pain, and an overall well-maintained general condition.

This relatively slow progression, in the absence of rapid visceral dissemination, suggests a low-grade or indolent carcinoma, compatible with certain differentiated urothelial tumor variants exhibiting an atypical immunohistochemical profile (notably loss of GATA3 expression).

A multidisciplinary management approach was undertaken, involving the urology, medical oncology, and pathology teams.

Locally, control of the inguinal mass relied on daily wound care, including cleansing of the cutaneous fistula, application of antiseptic dressings, and prevention of secondary infection. Empirical antibiotic therapy was initiated, resulting in local improvement and progressive drying of the wound.

Systemically, after multidisciplinary discussion, systemic chemotherapy was not initiated due to the high risk of toxicity and the expected limited tolerance in a 91-year-old patient.

An oral or carboplatin-based chemotherapy regimen was considered but ultimately not administered due to the patient's frailty and borderline renal function.

Immunotherapy, which is recommended for locally advanced or metastatic urothelial carcinoma in patients unfit for platinum-based regimens, was discussed but not initiated in the absence of a histologically confirmed primary tumor.

Radiological follow-up showed persistent necrotic left inguinal lymphadenopathies, with increased size and ongoing cutaneous

fistulization, but no evidence of new distant metastases.

Clinically, the patient remained stable for several months, with preserved diuresis, absence of pelvic pain, and overall good general condition.

This slow progression, without rapid visceral dissemination, supports the hypothesis of a low-grade or indolent carcinoma, consistent with certain differentiated urothelial tumor variants exhibiting an atypical immunohistochemical profile (loss of GATA3 expression).

Discussion

This case illustrates the diagnostic and therapeutic complexity posed by metastatic urothelial carcinoma (UC) tumors without an identified primary site (i.e., "urothelial carcinoma of occult origin"). Although urothelial origin is the most common tumor type in the genitourinary system, presentation with fistulized inguinal lymphadenopathy in an elderly patient, in the absence of a bladder or ureteral tumor visible on imaging and endoscopy, remains an exceptional entity.

Carcinoma of Unknown Primary (CUP) represents approximately 3-5% of malignant solid tumors and is associated with an often poor prognosis due to delayed diagnosis and lack of targeted treatment [9]. In this context, UC of occult origin is even rarer, and few series report it [10]. The fact that our patient presented with a pelvic and inguinal lymph node mass, suspicious ureteral thickening, and a bladder diverticulum with parietal irregularity reinforces the hypothesis of a urothelial origin, although this has not been confirmed.

In the literature, a few isolated cases describe metastases (pulmonary, lymph node, bone) revealing occult UC, such as the case reported by Bu et al. (2019) in a patient with multiorgan involvement without an identifiable primary lesion [10]. Our case is consistent with this and illustrates one of the most extreme forms of the diagnostic challenge in urology-oncology.

The diagnosis of occult UC requires a comprehensive exploration using morphological imaging: thoracoabdominopelvic CT, MRI if necessary, and PET-CT to identify occult foci. In our case, the TAP revealed enhanced ureteral thickening (36 mm) and a right lateral bladder diverticulum, two structural clues that could suggest a small or infiltrating primary site.

Cystoscopy/ureteroscopy is essential to visualize any macroscopic bladder or ureteral lesion. Here, the cystoscopy was normal, except for the diverticulum without a visible lesion. Targeted ureteroscopy of the ureteral thickening area could have added additional information. Molecular biology using NGS panels or molecular signatures (basal vs. luminal classifications, genomic alterations such as FGFR3, ERCC2, TP53) could help confirm the urothelial origin and guide treatment (e.g., targeted therapy) or follow-up strategy, especially in the absence of a visible primary site.

The use of an immunohistochemical panel is essential to guide tumor origin in cases of isolated metastasis. The GATA3 marker is widely used in pathology to identify urothelial differentiation [11]. However, several studies have shown that loss of GATA3 expression can occur in high-grade UC and dedifferentiated forms, limiting its diagnostic value in isolation.

For example, in a study of 2,710 bladder carcinomas, GATA3 positivity was 59.9% for stages pT2-4, compared to 98-99% for superficial tumors (pTa) [12]. This loss of expression has been correlated with increased tumor aggressiveness and greater invasiveness/metastasis [13]. The study by Li et al. has also shown that GATA3 deletion induces increased migration and invasion of urothelial cancer cells via activation of MMP-2 and MMP-9, suggesting a suppressive role for GATA3 in the urothelium [14].

In this case, the CK7+/CK20+, p63+, p40+, GATA3–immunophenotype is consistent with a dedifferentiated entity and suggests that GATA3 loss should not rule out the possibility of urothelial UC. Furthermore, some variants (plasmacytoid, sarcomatoid) frequently exhibit GATA3 absence [15]. It is therefore recommended to use an extensive panel (uropap, CK5/6, p40/p63, GATA3, etc.) and to correlate with clinical and imaging findings.

Typically, urothelial bladder carcinomas spread via the lymphatic system to the obturator, internal/external iliac, and then common iliac lymph nodes. A primary inguinal location is very rare [16]. The fact that our case presented with large left inguinal lymphadenopathy and an ulcerative-necrotizing mass invading the femoral region suggests atypical lymph node extension or secondary dissemination through unusual collateral drainage, possibly related to an undetected primary ureteral or lower bladder lesion.

Therapeutic choice is all the more complex in this type of case because the absence of an identified primary site limits the application of standardized protocols for urothelial cancer. Treatments such as platinum-based chemotherapy or anti-PD-1/PD-L1 immunotherapy are generally recommended for metastatic urothelial UC. However, in a 91-year-old patient with borderline renal function.

Conclusion

This case highlights the rarity and diagnostic challenge of occult primary urothelial carcinoma, which may initially present as isolated inguinal lymphadenopathy without detectable bladder or ureteral involvement. The integration of morphological, radiological, and immunohistochemical data remains essential to guide diagnosis in such atypical clinical presentations.

Although GATA3 negativity is uncommon, it does not exclude a urothelial origin, particularly in poorly differentiated or dedifferentiated forms. This case also underscores the importance of a multidisciplinary approach involving urology, pathology, and oncology teams to develop an appropriate diagnostic and therapeutic strategy tailored to the patient's condition.

In elderly and frail patients, management should focus on symptom control, infection prevention, and close follow-up. Finally, such rare presentations emphasize the need for further documentation to improve understanding of the atypical lymphatic dissemination mechanisms of urothelial carcinoma.

References

1. Le Chevalier T, Cvitkovic E, Caille P (1988) Early metastatic cancer of unknown primary origin at presentation. A clinical study of 302 consecutive autopsied patients. *Arch Intern Med* 148: 2035–2039.
2. Desai VB, Wright JD, Schwartz PE (2018) Occult gynecologic cancer in women undergoing hysterectomy or myomectomy for benign indications. *Obstet Gynecol* 131: 642–651.

3. Bond LM, Skrobo D (2015) Multiple embolic cerebral infarcts as the first manifestation of metastatic ovarian cancer. *BMJ Case Rep* 2015: bcr2015211521.
4. Tsai CC, Wu MN (2015) Frequent ischemic stroke as first manifestation of occult colon cancer: a rare case. *Am J Case Rep* 16: 723–727.
5. Galimberti V, Bassani G, Monti S (2004) Clinical experience with axillary presentation breast cancer. *Breast Cancer Res Treat* 88: 43–47.
6. Walker GV, Smith GL, Perkins GH (2010) Population-based analysis of occult primary breast cancer with axillary lymph node metastasis. *Cancer* 116: 4000–4006.
7. Pavlidis N, Pentheroudakis G (2012) Cancer of unknown primary site: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 23: 86–91.
8. Bu K (2019) An occult urothelial carcinoma with wide multiorgan metastasis. *Med (Baltimore)* 98: e15709.
9. Chu PG, Weiss LM (2014) GATA3: a multi-specific marker with emerging utility in surgical pathology. *Mod Pathol* 27: 19–30.
10. Plage H (2022) GATA3 expression loss is linked to stage progression but is unrelated to prognosis in muscle-invasive urothelial carcinoma of the bladder. *Mod Pathol* 35: 1234–1244.
11. Li Y, Ishiguro H (2014) Loss of GATA3 in bladder cancer promotes cell migration and invasion. *Cancer Biol Ther* 15: 428–435.
12. Isaka M (2021) Diagnostic relevance of GATA3 expression in urinary bladder carcinoma of divergent differentiation and other histological variants. *Indian J Pathol Microbiol* 64: 560–566.
13. Miettinen M, McCue PA (2014) GATA3: a multispecific but potentially useful marker in surgical pathology—a systematic analysis of 2500 epithelial and non-epithelial tumors. *Am J Surg Pathol* 8: 13–22.
14. Leissner J (2004) Anatomic extent of pelvic lymph node dissection: impact on outcome in patients with bladder cancer. *Eur Urol* 45: 242–249.
15. Powles T, van der Heijden MS, Castellano D (2018) Atezolizumab versus chemotherapy in patients with platinum-treated locally advanced or metastatic urothelial carcinoma (IMvigor211): a multicentre, open-label, phase 3 randomised controlled trial. *Lancet* 391: 748–757.
16. Bellmunt J, de Wit R, Vaughn DJ (2017) Pembrolizumab as second-line therapy for advanced urothelial carcinoma. *N Engl J Med* 376: 1015–1026.

Copyright: ©2025 Bjane O, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.