

Closing the Waste Loop: Recycling, Resource Recovery and the Environmental Health Consequences of Poor Waste Management

Satish Bhagwatrao Aher* and Pooja Saroj

Division of Environmental Monitoring and Exposure Assessment (Air), ICMR National Institute for Research in Environmental Health, Bhopal 462030, Madhya Pradesh, India

ABSTRACT

Rapid urbanization, industrialization, changing consumption patterns and population growth have substantially increased the generation and diversity of solid waste. When waste is inadequately segregated, recycled or treated, it becomes an important source of environmental pollution through particulate matter, gaseous pollutants, greenhouse gases, toxic metals, hazardous chemicals, leachate and microbial bioaerosols. At the same time, a considerable proportion of discarded materials retains significant potential for recycling, material recovery, composting, biomethanation and energy recovery. Consequently, waste management represents both an environmental health challenge and an opportunity for resource conservation and pollution prevention. This review synthesizes evidence on waste recycling, resource recovery and pollution generation across municipal, industrial, metallurgical, mining and small-scale recycling systems, with particular emphasis on the environmental-health consequences of inadequate waste management. The reviewed evidence demonstrates that open dumping, uncontrolled burning and poorly managed landfills are associated with particulate pollution, toxic emissions, methane generation, leachate contamination, heavy-metal exposure and microbial bioaerosols. In contrast, source segregation, material recovery, recycling, organic-waste treatment, landfill-gas recovery and appropriate industrial-waste utilization can divert recoverable materials from disposal pathways and potentially reduce environmental emissions. Quantitative evidence from recycling scenarios indicates that increasing recycling rates can substantially enhance energy savings and reduce modelled emissions, although the magnitude of benefits depends on material type, technology, collection systems and the environmental performance of recycling processes. The review also highlights an important paradox: recycling itself may generate occupational and environmental exposures when carried out in poorly controlled informal or small-scale facilities. A health-oriented circular waste-management approach therefore requires not only increasing recycling rates but also ensuring safe collection, segregation, processing and residual disposal. Future waste-management strategies should integrate material recovery, pollution control, occupational protection, environmental monitoring, life-cycle assessment and environmental justice. Closing the waste loop should consequently be viewed not simply as a resource-efficiency strategy but as an essential component of environmental health protection.

*Corresponding author

Satish Bhagwatrao Aher, Division of Environmental Monitoring and Exposure Assessment (Air), ICMR National Institute for Research in Environmental Health, Bhopal 462030, Madhya Pradesh, India.

Received: August 21, 2026; **Accepted:** September 01, 2026; **Published:** September 15, 2026

Keywords: Waste Recycling, Resource Recovery, Environmental Health, Waste Pollution, Particulate Matter, Heavy Metals, Bioaerosols, Landfill, Informal Recycling, Circular Economy

Introduction

Waste is an unavoidable consequence of human consumption, production and economic activity [1]. However, waste becomes an environmental-health problem when materials that retain economic or resource value are mixed, discarded, openly burned or deposited in poorly managed disposal sites [2]. The environmental consequences extend beyond the physical accumulation of waste and include deterioration of air, water and soil quality, greenhouse-gas emissions, occupational exposure and risks to communities living near waste-management facilities [3-5]. Previous reviews have emphasized that the waste sector is closely connected with atmospheric pollution and climate change through landfill emissions, open burning, uncontrolled decomposition and inefficient treatment systems [6,7]. The problem is particularly important in rapidly urbanizing developing regions, where waste-generation rates are increasing while collection, segregation, recycling and

scientific treatment systems frequently remain inadequate. In India, for example, the reviewed literature reports substantial municipal solid-waste generation accompanied by persistent gaps in source segregation, material recovery and scientific residual disposal [8-10]. Open dumping remains an important pathway through which waste is transformed from a potentially recoverable resource into a source of environmental contamination. Recent synthesis of Indian dumping sites has identified particulate matter, sulfur dioxide, nitrogen dioxide, carbon monoxide, methane, volatile organic compounds, bioaerosols and odour among the major pollution concerns associated with unmanaged waste disposal [3].

The environmental burden of waste is therefore closely linked to the extent to which materials are recovered before disposal. Recycling can conserve raw materials, reduce energy requirements and divert materials from landfills and dumpsites. A case study from Shiraz, Iran, demonstrated this potential quantitatively: increasing the recycling rate from 15% toward higher recycling scenarios was associated with substantial increases in energy savings and reductions in modelled pollutant emissions Farzadkia

et al. [11]. Similarly, studies of industrial and metallurgical wastes have identified opportunities for recovering metals and utilizing mineral residues in construction materials, thereby reducing the quantity of waste requiring long-term storage or disposal [13,14]. However, recycling should not automatically be considered environmentally benign. Small-scale and informal recycling operations may generate toxic fumes, contaminated wastewater, metal-containing residues and smoke when appropriate pollution-control technologies are absent. This issue is particularly important for e-waste, batteries, metals and plastics, where uncontrolled processing can transfer contaminants from solid waste into air, water and soil [15]. Waste pickers and workers involved in collection and sorting may also experience disproportionate exposure to particulate matter, hazardous gases, microbial agents and other occupational hazards [16].

Thus, the central challenge is not simply to recycle a larger proportion of waste. The objective should be to establish a safe recycling and resource-recovery system that simultaneously reduces waste disposal, conserves resources and minimizes environmental-health risks. This review examines this interconnected relationship by synthesizing evidence on waste recycling, resource recovery, pollution pathways and environmental-health implications. Particular attention is given to municipal solid waste, industrial and metallurgical residues, mining waste, informal recycling and landfill resource recovery. The review further identifies barriers and proposes a health-oriented framework for closing the waste loop.

Approach and Evidence Synthesis

This review synthesizes published studies compiled in the evidence table developed for the present analysis (Table 1-3). The evidence base comprises studies covering municipal solid waste, industrial and metallurgical residues, mining waste, recycling industries, landfills and waste-to-energy systems from India and other countries. The studies were examined according to five interconnected dimensions: (i) waste type and management system, (ii) recycling or resource-recovery pathway, (iii) pollution generated, (iv) environmental and health implications, and (v) opportunities and limitations associated with improved waste recovery. The evidence includes both quantitative studies and broader reviews. Importantly, the available literature does not consistently report recycling quantities or recovery efficiencies. Several studies focus primarily on pollution from disposal sites and therefore provide evidence of the environmental burden associated with failure to recover materials rather than direct measurements of recycling performance. Consequently, recycling rates reported in individual studies are interpreted as study-specific values or scenario estimates and are not treated as directly comparable national indicators. This distinction is important because waste recycling, material recovery, energy recovery, composting, bi-methanation and landfill-gas recovery represent different processes and cannot be treated as equivalent measures of recycling. The present synthesis therefore focuses on the broader concept of resource recovery, while distinguishing material recycling from energy recovery wherever the source information permits.

Waste Generation, Disposal and the Environmental-Health Burden

Municipal Solid Waste and the Transition from Resource to Pollutant

Municipal solid waste contains a heterogeneous mixture of biodegradable materials, paper, cardboard, plastics, metals, glass, textiles and other materials. Many of these fractions have

direct recycling or recovery potential. Nevertheless, inadequate segregation frequently results in valuable materials being mixed with organic and hazardous fractions, reducing their recoverability and increasing the environmental burden of subsequent disposal. Indian waste-management reviews consistently identify source segregation, material recovery and scientific disposal as major weaknesses. Ahluwalia and Patel emphasized that collection and transportation often receive greater attention than segregation, recycling and scientific treatment [9]. Similarly, Gour & Singh identified inadequate segregation, limited recovery and enforcement gaps as recurring challenges in India's diverse waste streams [8]. When recoverable materials are instead disposed of at open dumps, several pollution pathways are activated simultaneously. Decomposition produces methane and carbon dioxide, while open burning produces particulate matter, carbon monoxide, nitrogen oxides, sulfur oxides, volatile organic compounds and toxic combustion products. Leachate may transport dissolved organic matter, metals and other contaminants into surrounding soil and water. Thus, the environmental cost of failing to recycle is not restricted to the loss of material value; it includes the creation of additional pollution burdens as in Figure 1.




Figure 1: Unsustainable Waste-Management Pathway and its Cumulative Environmental, Social, Economic, and Human-Health Impacts

Open Dumping and Uncontrolled Burning

Evidence from Indian dumping sites demonstrates the magnitude of pollution associated with poorly managed waste. Bamotra et al. reported high concentrations of PM_{2.5} and PM₁₀ across reviewed Indian dumping sites, together with gaseous pollutants, methane, volatile organic compounds, bioaerosols and odour [3]. Sethy et al. investigated the Bhuasuni/Vuasuni dumping site in Bhubaneswar and identified particulate matter as the primary contributor to deterioration in local air quality [16]. Similarly, Pathak & Kushwaha reported elevated suspended and respirable particulate matter around municipal dumping sites in Madhya Pradesh, with windblown dust, vehicle movement and open burning contributing to pollution [17]. The evidence from Ahmedabad provides an additional perspective on the toxic nature of waste-related particulate pollution. At the Pirana landfill, Mehta et al. reported PM_{5.0} concentrations of approximately 189–1,600 µg m⁻³, with particle-bound metals including Cu, Fe, Ni, Co, Al, As, Pb, Cr, Cd, Zn and Hg [18]. Solid-waste burning together with industrial coal combustion contributed to a measurable fraction of the metal pollution. These observations demonstrate that disposal sites can become localized sources of complex mixtures of particulate and chemical contaminants rather than merely sites of solid-waste accumulation.

Landfill Gases and Climate-Related Pollution

Organic waste deposited in landfills undergoes anaerobic decomposition and generates methane, carbon dioxide and trace gases. Methane is particularly important because it is both a greenhouse gas and a potentially recoverable energy resource. Mor et al. estimated methane generation of approximately 0.34 Gg year⁻¹ from the Chandigarh municipal solid-waste disposal system and demonstrated the potential for landfill-gas recovery to generate electricity [19]. This represents an important shift in the waste-management paradigm. Rather than treating landfill methane exclusively as a pollutant, methane can be viewed as an indicator of recoverable energy that becomes available because biodegradable material has entered a disposal pathway. Nevertheless, energy recovery from landfill gas should not be considered a substitute for source segregation and organic-waste diversion. Prevention and biological treatment of biodegradable waste remain preferable to allowing organic matter to decompose anaerobically in landfills.

Recycling and Resource Recovery as Pollution-Prevention Strategies

Municipal Recycling

Recycling can interrupt the pathway connecting waste generation with environmental pollution. When paper, cardboard, glass, metals and plastics are separated at source and directed toward appropriate recovery systems, the quantity of material requiring disposal decreases. The benefits include conservation of raw materials, energy savings and potential reductions in emissions associated with primary production. The quantitative evidence from Farzadkia et al. is particularly useful in demonstrating this relationship [11]. In Shiraz, the baseline recycling rate was approximately 15%, while scenarios involving substantially greater recycling were evaluated for major recyclable materials. The study estimated annual recovery of approximately 11,912 tonnes across the evaluated material categories under the reported scenario and showed that increasing recycling could substantially increase energy savings and reduce modelled emissions [11]. Although this result is location-specific, it illustrates a general principle: the environmental benefit of recycling is determined not only by the mass of waste recovered but also by the energy and emissions displaced in the production of virgin materials.

Table 1: Industrial, Metallurgical and Mining Waste: Resource Recovery and Pollution-Control Opportunities

Study / location	Waste stream	Pollution and resource-recovery evidence	Key environmental-health implication
Chhattisgarh, India [20]	Iron and steel waste: slag, flue dust, sludge and process residues	PM, SO ₂ , NO _x , CO, CO ₂ , CH ₄ , VOCs, PAHs and PCDD/Fs; potential heavy-metal leaching from slag. Bhilai generates ~2.1 million t slag/year; ~81% reported as blast-furnace air-cooled/granulated slag.	Slag utilization can reduce disposal burdens, but lined storage, leachate monitoring and emission controls are required to minimize environmental and health risks.
India [12]	Red mud, spent pot lining, fly ash, slags, dross, dust, sludge, tailings and leach residues	Dust, fluoride and metal fumes, with metal/fluoride/cyanide-bearing leachate and contaminated wastewater. ~2 million tons of red mud per year reported; selected residues have potential for metal recovery and use in cement, bricks and roads.	Resource recovery can reduce hazardous waste accumulation, but uncontrolled residues and informal lead recycling pose significant environmental-health risks.
Odisha, India [13]	Iron-ore tailings, overburden, slag, dust and mineral residues	Dust/PM, process emissions, metal-rich runoff/leachate and land degradation. Potential applications include bricks, concrete/cement, roads, backfilling and metal/mineral recovery.	Beneficial utilization can divert large volumes of mining waste from disposal; characterization, quality standards and site-specific assessment are essential.

Abbreviations: PM, particulate matter; VOCs, volatile organic compounds; PAHs, polycyclic aromatic hydrocarbons; PCDD/Fs, polychlorinated dibenzo-p-dioxins/dibenzofurans.

Industrial and Metallurgical Waste Recovery

Industrial wastes present a different but potentially substantial resource-recovery opportunity because many residues contain concentrated mineral or metallic fractions. The Indian non-ferrous metallurgical sector generates red mud, spent pot lining, fly ash, slags, dross, dust, sludge and other residues. Agrawal et al. documented recovery and utilization options for these materials, including metal recovery and incorporation into cement, bricks and road-related applications [12]. Similarly, steel production produces large quantities of slag and other process residues. Tiwari et al. highlighted substantial slag generation in central Indian steel plants and identified material utilization as an important component of resource efficiency [20]. However, the environmental benefit of slag utilization depends on appropriate characterization, storage and processing because improperly stored residues may release metals and other contaminants to surrounding environments. Thus, industrial recycling should follow a “recover and control” principle: resource recovery should proceed only when the

recovered material can be safely processed and used without transferring contaminants from one environmental compartment to another.

Mining and Mineral Residues

Mining and mineral-processing wastes represent another large-volume resource stream. Iron-ore tailings, overburden, slag and dust can cause land degradation, particulate pollution and metal-rich runoff or leachate. Krishna et al. identified potential applications of steel and mining residues in bricks, concrete, cement, roads and backfilling, together with opportunities for mineral and metal recovery [13]. Because mining residues are generated in very large quantities, even partial utilization can substantially reduce land requirements for storage and disposal. However, material characterization, contaminant assessment and performance standards are essential before such residues are incorporated into construction materials.

E-waste and Small-Scale Recycling

Small-scale recycling represents one of the clearest examples of the dual nature of recycling. Recovery of metals, plastics and other components can conserve valuable resources, but poorly controlled processing may create significant pollution. Kojima & Jain reported pollution concerns associated with small-scale recycling of e-waste, printed circuit boards, lead batteries, metal scrap, paper and plastics [14]. Potential pollution pathways include toxic fumes, contaminated wastewater, metal-containing residues and smoke from uncontrolled plastic combustion. Therefore, an increase in recycling rate does not necessarily correspond to an equivalent improvement in environmental health. A recycling system that transfers hazardous substances from discarded products into uncontrolled emissions or wastewater may simply relocate the environmental burden.

Pollution Pathways Associated with Waste Management Air Pollution

Air pollution represents one of the most immediate environmental-health pathways associated with waste management. Particulate matter may arise from waste handling, vehicle movement, windblown dust, mechanical processing, open burning and industrial recycling. Combustion can additionally release CO, NO_x, SO_x, VOCs, PAHs, dioxins/furans and other hazardous pollutants. Waste-to-energy technologies also require careful assessment. Karmakar et al. reviewed Indian waste-to-energy technologies and highlighted emissions of particulate matter, nitrogen oxides, sulfur oxides, acid gases, VOCs, greenhouse gases, PAHs and persistent organic pollutants [10]. The review also emphasized that India's high moisture content, inadequate source segregation and relatively low calorific value can constrain waste-to-energy performance. Consequently, waste-to-energy should not be treated as an automatic alternative to recycling.

Water and Soil Contamination

Waste disposal and poorly controlled recycling can contaminate soil and water through leachate, wastewater and residue deposition. Heavy metals are particularly important in metallurgical waste, e-waste and mixed waste. Lead, cadmium, chromium, mercury, arsenic, nickel and other metals may persist in environmental media and can enter food chains or groundwater under suitable conditions. The environmental objective of recycling should therefore be to prevent contaminant release at source. Merely moving contaminated material from a landfill to an inadequately controlled recycling facility does not constitute sustainable resource recovery.

Greenhouse Gases

Waste management contributes to climate change primarily through methane generation from landfilled organic matter, carbon dioxide from waste combustion and energy use, and emissions associated with the production of virgin materials that could otherwise be displaced through recycling. Sharma et al. emphasized the relationship between improper waste management, atmospheric pollution and climate change [7]. Recycling and organic-waste diversion can therefore provide climate benefits, while landfill-gas capture offers a secondary recovery pathway for waste that has already entered the landfill system. Mor et al. demonstrated this potential in the context of Chandigarh [19].

Bioaerosols and Microbial Hazards

An important but comparatively less visible dimension of waste-related pollution is biological contamination. Landfills contain organic material that can support microbial growth and aerosolization. Srivastava et al. identified fungal bioaerosols including *Penicillium*, *Aspergillus*, *Rhizopus*, *Alternaria*, *Fusarium* and *Cladosporium* around the Okhla landfill in Delhi [21]. Madhwal et al. reported substantial bacterial and fungal bioaerosol concentrations at the Sahastradhara landfill in Dehradun, together with particulate matter and potentially pathogenic microorganisms [22]. The study also indicated that waste handling, meteorological conditions and biowaste burning influenced exposure patterns. These findings broaden the concept of waste pollution beyond conventional chemical contaminants and indicate that waste-management systems must also address biological exposure.

Environmental and Occupational Health Implications

The health consequences of waste pollution arise through multiple exposure pathways. Communities living near disposal sites may be exposed through inhalation of particulate matter and gases, ingestion of contaminated water or food, and dermal contact with contaminated soil or waste. Workers and waste pickers may experience much higher exposure because of direct handling, sorting, transportation and processing. The evidence synthesized by Damayanty et al. indicates that landfill workers, scavengers and nearby marginalized communities may experience exposure to H₂S, NH₃, VOCs, PM_{2.5}, PM₁₀, methane, carbon dioxide and microbial bioaerosols [15]. Reported health concerns include respiratory, infectious and cardiovascular effects, with possible carcinogenic risks associated with some exposures. This occupational dimension is critical for recycling policy. Waste pickers often perform an economically valuable material-recovery function, but their contribution can remain invisible in formal waste-management statistics. Formalization, personal protective equipment, safer sorting facilities, training, vaccination where appropriate, health surveillance and access to social protection should therefore be integrated into recycling strategies.

Table 2: Municipal Solid Waste Management, Recovery and Environmental-Health Implications

Study location (Reference)	Waste stream / management system	Pollution and recovery evidence	Key environmental-health implication
India [10]	MSW; incineration, gasification, pyrolysis and biochemical WtE	PM, NO _x , SO _x , CO, VOCs, GHGs and PCDD/Fs. India generated 160,038.9 TPD MSW; reported WtE efficiencies varied by technology.	High moisture, poor segregation and low calorific value may increase emission risks; source segregation and emission controls are essential.
India, national/city-level [9]	Biodegradable and dry recyclables; composting, biomethanation, material recovery and WtE	MSW generation estimated at 64–72 million t yr ⁻¹ , informal recovery is substantial but below potential. Open dumping generates methane, smoke and leachate.	Inadequate segregation and scientific disposal increase pollution. Decentralized processing and integration of waste pickers are recommended.
Multiple Indian dumpsites [3]	Mixed MSW at open dumps/landfills	PM _{2.5} , PM ₁₀ , SO ₂ , NO ₂ , CO, CH ₄ , VOCs, bioaerosols and odour. Reported ~5% recycling, 18% composting and 77% open dumping.	Dumpsites create combined climate, toxic-exposure and microbial risks; stronger mitigation and epidemiological evidence are needed.
Bhubaneswar, India [16]	Mixed MSW; open dumping/landfill	PM ₁₀ , PM _{2.5} , SO ₂ , NO _x , CO, dust and odour; AQI = 51.9. Recycling throughput not reported.	PM was the major air-quality concern. Dust suppression, vegetation and improved site management are required.
Madhya Pradesh, India [17]	Mixed MSW; open dumping	SPM/RSPM frequently exceeded limits; windblown dust, vehicle exhaust and open burning were major sources.	Poorly designed dumpsites increase particulate exposure; open burning and windblown emissions should be controlled.
India [8]	Municipal, industrial, biomedical, construction and household wastes	Dust, smoke, odour, GHGs and leachate; inadequate segregation and recovery highlighted.	India's waste system requires improved segregation, recovery, data, enforcement and integrated treatment.
India [6]	Waste-related open burning and landfill/dumpsite emissions	PM, CO, NO _x , SO ₂ , VOCs and toxic smoke; recycling was not evaluated.	Waste burning is an important urban emission source requiring improved inventories and air-quality management.
Chennai, India [23]	Fresh/aged MSW containing plastics, paper, metals, organics and textiles	Fresh waste contained ~30 wt% recyclables and ~62 wt% combustibles; CH ₄ , CO ₂ , odour, leachate and metals are concerns.	Significant resource-recovery potential exists; biomining requires waste characterization and pollution control.
Global [7]	MSW/wastewater; landfill, incineration, composting, biomethanation and recycling	CH ₄ , CO ₂ , PM, VOCs, odour, PCDD/Fs and leachate. Literature estimate: ~20% MSW recycled and 14% converted to energy.	Integrated 4R systems, organic-waste diversion and methane recovery can reduce pollution and climate impacts.
China [24]	MSW landfill, composting, incineration and informal recovery	Recyclables estimated at 8–10%; 2010 treatment: 79% landfill, 19% incineration and 2% composting. Emissions include CH ₄ , CO ₂ , PM, metals and PCDD/Fs.	Treatment choice strongly influences emissions; recycling, landfill-gas capture and emission controls are important.
Chandigarh, India [19].	MSW disposal; landfill-gas capture/WtE	~350 TPD waste generation/disposal; >50% organic matter; estimated landfill-gas recovery potential 0.27–0.77 MW.	High organic content promotes methane generation; gas recovery can reduce emissions and recover energy.
Ahmedabad, India [18]	Mixed MSW landfill with open burning	PM _{5.0} ~189–1,600 µg m ⁻³ with particle-bound Cu, Fe, Ni, Co, As, Pb, Cr, Cd, Zn and Hg.	Waste/coal combustion contributes to metal-rich particulate pollution and associated health risks.

Abbreviations: WtE, waste-to-energy; MSW, municipal solid waste; PM, particulate matter; VOCs, volatile organic compounds; GHGs, greenhouse gases; PCDD/Fs, polychlorinated dibenzo-p-dioxins/dibenzofurans; AQI, air quality index; SPM, suspended particulate matter; RSPM, respirable suspended particulate matter. **Note:** Reported quantities and percentages are study-specific and should not be interpreted as directly comparable recycling rates. “Recovery” includes material, biological and energy-recovery pathways where reported.

Informal Recycling and Environmental Justice

The environmental-health burden of waste is not distributed equally. Communities living close to dumpsites and landfills and workers engaged in informal collection and recycling can experience disproportionately high exposure. This creates an environmental-justice dimension to waste management. Informal recycling systems can nevertheless provide substantial material-recovery services. The appropriate response is therefore not simply to eliminate informal recovery activities but to integrate workers into safer formal systems wherever feasible. Such integration could combine the material-recovery knowledge and collection networks of informal workers with improved infrastructure, occupational safety and regulatory oversight. This approach is particularly important in developing countries, where informal recovery may remain an important component of actual material recycling even when official statistics do not fully capture it.

Table 3: Recycling Systems, Informal Recovery and Environmental-Health Risks

Study location (Reference)	Waste stream / recycling system	Pollution, exposure and recovery evidence	Key environmental-health implication
Global [25]	Municipal, industrial, agricultural, radioactive and hazardous waste	Air, water and soil pollution, GHGs, radiation and ecosystem impacts; quantitative recycling data not reported.	Waste prevention, recycling, controlled treatment and coordinated regulation should be integrated to reduce environmental and health impacts.
Shiraz, Iran [11]	Dry recyclables: glass, paper/cardboard, metals, PET and plastics	Current recycling ~15%; recycling scenario ranging from 50–80%, increased material recovery and modelled emission/energy benefits; ~11,912 t yr ⁻¹ recovered under the reported scenario.	Increased recycling can substantially reduce emissions and energy demand, demonstrating its pollution-prevention potential.
India and Japan [14]	E-waste/PCBs, lead batteries, metal scrap, paper and plastics	Heavy metals, fumes, toxic smoke, wastewater and residues; quantitative recycling rate not reported.	Uncontrolled small-scale recycling can create significant exposure risks; formalization, pollution controls and enforceable standards are needed.
Global [26]	Mixed municipal and industrial waste	Air, water and soil pollution, GHGs and hazardous chemical/radiation impacts; quantitative recovery not reported.	Waste prevention and integrated management are preferable to disposal-focused approaches for protecting ecosystems and health.
Global [15]	Landfills and informal waste picking/scavenging	H ₂ S, NH ₃ , VOCs, PM _{2.5} , PM ₁₀ , CH ₄ , CO ₂ and microbial bioaerosols; waste pickers contribute to recyclable recovery.	Informal workers and nearby communities may experience disproportionate exposures, highlighting the need for occupational protection, biomonitoring and social inclusion.
Delhi, India [21]	MSW dump; waste handling and fungal bioaerosols	Fungal bioaerosols including <i>Penicillium</i> , <i>Aspergillus</i> , <i>Rhizopus</i> , <i>Alternaria</i> , <i>Fusarium</i> and <i>Cladosporium</i> ; site capacity ~2,000 t day ⁻¹ .	Respirable bioaerosols represent an important occupational and community exposure pathway, influenced by season and meteorology.
Dehradun, India [22]	Open MSW landfill; bacterial/fungal bioaerosols and PM	Fungal bioaerosols: 4,582.75 ± 1,358.25 CFU m ⁻³ , bacterial bioaerosols: 3,609.53 ± 1,000.28 CFU m ⁻³ , ~200 t day ⁻¹ dumped.	Waste handling and biowaste burning contribute to PM and bioaerosol exposure, with potential respiratory risks for workers and nearby populations.

Abbreviations: PM, particulate matter; VOCs, volatile organic compounds; GHGs, greenhouse gases; PET, polyethylene terephthalate; PCBs, printed circuit boards; CFU, colony-forming units.

Barriers to Safe and Sustainable Recycling

Several interrelated barriers limit the environmental, economic and health benefits that can be achieved through waste recycling and resource recovery. Inadequate source segregation is a fundamental constraint because mixing waste streams reduces the quality and recoverability of recyclable materials while increasing contamination of organic fractions. Effective recycling therefore depends on segregation at the point of generation. This challenge is compounded by inadequate recycling infrastructure, including insufficient collection systems, sorting facilities, material-recovery infrastructure and reliable markets for recovered materials, which can result in otherwise recyclable materials ultimately reaching dumpsites or landfills. At the same time, recycling processes themselves can become sources of pollution when air emissions, wastewater, leachate and hazardous residues are poorly controlled, with particular concerns for the recycling of metals, e-waste,

batteries and plastics. Weak regulatory enforcement further exacerbates these problems, as the existence of regulations does not necessarily ensure their implementation; effective scientific disposal, emission control, lined storage, leachate management and environmental monitoring require consistent enforcement and institutional capacity. Another important limitation is the lack of reliable and standardized data on actual recycling and recovery. Many studies report waste generation or disposal quantities without providing corresponding information on the amount actually recovered, making it difficult to compare recycling performance across cities, sectors and countries and to evaluate the effectiveness of waste-management interventions. Finally, informal-sector vulnerability represents an important social and occupational dimension of recycling. Informal workers may make substantial contributions to resource recovery but often operate without adequate occupational protection, health safeguards or

social security. Their formalization, inclusion in organized recycling systems and provision of appropriate occupational protection are therefore essential for developing a recycling system that is not only resource-efficient but also environmentally safe, socially equitable and protective of human health. These interconnected barriers and their consequences are summarized in Figure 2.



Figure 2: Interconnected Barriers Limiting Safe and Sustainable Recycling

A Health-Oriented Framework for Closing the Waste Loop

The evidence synthesized in this review supports a shift from a conventional “collect–transport–dump” model toward a “segregate–recover–treat–protect–monitor” framework. The first stage should be waste prevention and source segregation. The second should prioritize material recovery and recycling of paper, metals, glass, plastics and other technically recoverable materials. Biodegradable waste should be directed toward composting, anaerobic digestion or other suitable biological treatment systems. Industrial and mineral residues should undergo characterization before beneficial utilization or recovery. Only residual waste that cannot reasonably be recovered should proceed to controlled thermal treatment or engineered landfill disposal. Where landfilling is unavoidable, methane capture, leachate management, groundwater monitoring and air-quality surveillance should be implemented. The final component should be environmental-health protection. Air pollutants, heavy metals, hazardous chemicals, bioaerosols and occupational exposures should be monitored throughout the waste-management chain. Thus, recycling performance should be evaluated using not only “how much waste is recycled?” but also “how safely is it recycled?”

A useful conceptual relationship can be expressed as:

Waste generation → segregation → material/organic recovery → controlled processing → reduced disposal → reduced pollution → reduced exposure → improved environmental health

Conversely:

Mixed waste → dumping/burning → air/water/soil pollution → occupational/community exposure → environmental-health burden




Figure 3: Sustainable Waste-Management Pathway Illustrating a Health-Oriented Framework for Closing the Waste Loop through Source Segregation, Material and Organic Recovery, Industrial and Mineral Resource Recovery, and Controlled Treatment of Residual Waste, with Associated Environmental and Human-Health Benefits

This framework emphasizes that recycling and environmental health are not separate policy domains. They are interconnected components of a single waste-management system as shown in Figure 3.

Research Gaps and Future Priorities

The evidence base identifies several important research priorities. First, standardized reporting of waste generation, collection, segregation, recycling, recovery and final disposal is required. Without harmonized indicators, comparisons among cities and countries remain difficult. Second, future studies should quantify the environmental benefits of recycling using life-cycle assessment, material-flow analysis and emission inventories. Recycling should be compared with landfilling, incineration and other treatment options using consistent system boundaries. Third, greater attention is required to emissions from recycling facilities themselves. The environmental-health literature has extensively documented pollution from dumpsites, but comparatively fewer studies quantify emissions and exposures from formal and informal recycling operations. Fourth, occupational-health studies should move beyond short-term cross-sectional assessments toward longitudinal studies, biomonitoring and exposure assessment among waste workers and informal recyclers. Damayanty et al. specifically highlighted the need for stronger longitudinal evidence [15]. Fifth, emerging waste streams-including electronic waste, plastics, batteries, photovoltaic materials and composite materials-require integrated assessment of resource recovery potential and environmental-health risks. Finally, recycling policy should be evaluated using a broader set of indicators that combine resource recovery, pollution reduction, greenhouse-gas mitigation, occupational safety, community exposure and environmental justice.

Conclusions

Waste recycling and resource recovery offer major opportunities to reduce the environmental burden associated with waste disposal while conserving materials and energy. However, the reviewed evidence demonstrates that the benefits of recycling cannot be assessed solely by recycling rates. Poorly managed waste disposal produces particulate matter, toxic gases, greenhouse gases, heavy metals, leachate and microbial bioaerosols, while poorly controlled recycling can itself generate hazardous emissions and occupational exposures. The most effective approach is therefore an integrated waste-management hierarchy that prioritizes prevention, source segregation, material recovery, biological treatment of organic waste, safe industrial residue utilization and controlled treatment of unavoidable residuals. Landfill-gas recovery and waste-to-energy can contribute to resource recovery under appropriate conditions but should not substitute for source reduction and material recycling. A transition toward a circular waste-management system should consequently be accompanied by strong environmental-health safeguards. Recycling facilities must control air emissions, wastewater and hazardous residues; workers must receive appropriate occupational protection; informal recyclers should be integrated into safer systems; and communities surrounding waste facilities should benefit from environmental monitoring and transparent risk communication. Ultimately, closing the waste loop should mean more than converting waste into a usable product. It should mean preventing the conversion of recoverable resources into pollutants and protecting workers, communities and ecosystems throughout the waste-management chain. A health-oriented approach to recycling can therefore serve simultaneously as a strategy for resource conservation, pollution prevention, climate mitigation and environmental-health protection.

Acknowledgements

The authors express their sincere gratitude to the Director, ICMR National Institute for Research in Environmental Health, Bhopal for providing guidance and support.

Funding

The authors declare that no specific external funding was received for conducting this study.

Conflict of Interest

The authors declare that there are no conflicts of interest regarding the publication of this manuscript.

Author Contributions (CRediT Statement)

Satish Bhagwatrao Aher: Conceptualization, Methodology, Graphics, Writing – original draft, Writing – review & editing.

Pooja Saroj: Literature survey, Graphics, Writing – original draft, Writing – review & editing.

All authors have read and approved the final version of the manuscript.

Data Availability Statement

Not applicable

Ethical Approval

Not applicable

Informed Consent

Not applicable

Consent for Publication

All authors have reviewed and approved the manuscript for publication.

References

1. Amasuomo E, Baird J (2016) The concept of waste and waste management. *J Mgmt & Sustainability* 6: 88.
2. Ayomoh M K O, Oke S A, Adededeji W O, Charles-Owaba O E (2008) An approach to tackling the environmental and health impacts of municipal solid waste disposal in developing countries. *Journal of Environmental Management* 88: 108-114.
3. Bamotra S, Yadav S, Tandon A (2025) Air pollution from dumping sites in India: implications on environmental health. *Water Air & Soil Pollution* 236: 624.
4. Bisht A, Kamboj V, Kamboj N, Bharti M, Bahukahndi K D (2024) Impact of solid waste dumping on soil quality and its potential risk on human health and environment. *Environmental Monitoring and Assessment* 196: 763.
5. Siddiqua A, Hahladakis J N, Al-Attiya W A K (2022) An overview of the environmental pollution and health effects associated with waste landfilling and open dumping. *Environmental Science and Pollution Research* 29: 58514-58536.
6. Guttikunda S K, Goel R, Pant P (2014) Nature of air pollution, emission sources, and management in the Indian cities. *Atmospheric environment* 95: 501-510.
7. Sharma B, Vaish B, Srivastava V, Singh S, Singh P, et al. (2018) An Insight to Atmospheric Pollution- Improper Waste Management and Climate Change Nexus. In: Oves, M., Zain Khan, M., M.I. Ismail, I. (eds) *Modern Age Environmental Problems and their Remediation*. Springer, Cham.
8. Gour A A, Singh S K, Mandal A (2023) Solid waste management in India: a state-of-the-art review. *Environmental Engineering Research* 28: 220249.
9. Ahluwalia I J, Patel U (2018) Solid waste management in India: An assessment of resource recovery and environmental impact (No. 356). Working paper 1-48.

10. Karmakar A, Daftari T, Chandan M R, Shaik A H, Kiran B, Chakraborty S (2023) A comprehensive insight into Waste to Energy conversion strategies in India and its associated air pollution hazard. *Environmental Technology & Innovation* 29: 103017.
11. Farzadkia M, Mahvi A H, Norouzian Baghani A, Sorooshian A, Delikhoon M, et al. (2021) Municipal solid waste recycling: Impacts on energy savings and air pollution. *Journal of the Air & Waste Management Association* 71: 737-753.
12. Agrawal A, Sahu KK, Pandey BD (2004) Solid waste management in non-ferrous industries in India. *Resources Conservation and Recycling* 42: 99-120.
13. Krishna R S, Mishra J, Meher S, Das S K, Mustakim S M, et al. (2020). Industrial solid waste management through sustainable green technology: Case study insights from steel and mining industry in Keonjhar, India. *Materials today: proceedings* 33: 5243-5249.
14. Kojima M, Jain A (2008) Controlling Pollution in Small-Scale Recycling Industries: Experiences in India and Japan. In M. Kojima (Ed.), *Promoting 3Rs in Developing Countries: Lessons from the Japanese Experience*. Institute of Developing Economies 36-58.
15. Damayanty S, Saleh L M, Mallongi A, Noor N N, Daud A, et al. (2025) Airborne pollutants from landfills and their health effects on scavengers and nearby communities: A scoping review. *Edelweiss Applied Science and Technology* 9: 1631-1651.
16. Sethy K M, Harichandan A, Pradhan S, Ghosh M, Patra S H (2019) Threat to ambient air quality due to poor municipal solid waste management: A case study in and around vuasuni dumping site, Bhubaneswar, Odisha, India. *Journal of Environment and Earth Science* 9: 7-16.
17. Pathak V, Kushwaha BP (2012) Study on ambient air quality at selected few municipal solid waste dumping sites of district Satna (M.P.) India. *Life Science Bulletin* 9: 39-42.
18. Mehta UH, Kaul DS, Westerdahl D, Ning Z, Zhang K, et al. (2022) Understanding the sources of heavy metal pollution in ambient air of neighboring a solid waste landfill site. *Aerosol Science and Engineering* 6: 161-175.
19. Mor S, Kaur K, Ravindra K (2024) Methane emissions from municipal landfills: a case study of Chandigarh and economic evaluation for waste-to-energy generation in India. *Frontiers in Sustainable Cities* 6: 1432995.
20. Tiwari MK, Bajpai S, Dewangan UK (2016) Air and leaching pollution scenario by iron and steel plants, in central India. *Elixir Pollution* 101: 44011-44017.
21. Srivastava A, Verma R, Mehta D (2021) Characterization of bioaerosols in and around a landfill site in Delhi. *Aerobiologia* 37: 585-596.
22. Madhwal S, Prabhu V, Sundriyal S, Shridhar V (2020) Distribution, characterization and health risk assessment of size fractionated bioaerosols at an open landfill site in Dehradun, India. *Atmospheric Pollution Research* 11: 156-169.
23. Peter AE, Nagendra SS, Nambi I M (2019) Environmental burden by an open dumpsite in urban India. *Waste management* 85: 151-163.
24. Tian H, Gao J, Hao J, Lu L, Zhu C, et al. (2013) Atmospheric pollution problems and control proposals associated with solid waste management in China: a review. *Journal of hazardous materials* 252: 142-154.
25. Ibro Skenderovic, Becir Kalac, Suad Becirovic (2015) Environmental pollution and waste management. *Balkan Journal of Health Science* 3: 2-10.
26. Budjav B (2022) Evaluation of Environmental Pollution and Waste Management Strategies on the Ecosystem. *Journal of Enterprise and Business Intelligence* 223-234.